

Document Version

Final published version

Citation (APA)

Hoogeman, M., Arts, T., van de Water, S., van der Voort, S., Perko, Z., Lathouwers, D., Breedveld, S., & Heijmen, B. (2016). SP-0010: Selection of patients for proton therapy: a physicists view. *Radiotherapy & Oncology*, 119(Suppl. 1), S4. [https://doi.org/10.1016/S0167-8140\(16\)31259-2](https://doi.org/10.1016/S0167-8140(16)31259-2)

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

In case the licence states "Dutch Copyright Act (Article 25fa)", this publication was made available Green Open Access via the TU Delft Institutional Repository pursuant to Dutch Copyright Act (Article 25fa, the Taverne amendment). This provision does not affect copyright ownership.
Unless copyright is transferred by contract or statute, it remains with the copyright holder.

Sharing and reuse

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

attention to the radiotherapy planning and delivery elements, and careful systematic and prospective documentation of tumor and normal tissue outcomes. Even if randomised trials are deemed unsuited to the setting, protocol based approaches in registered phase I/II trials are appropriate to enhance standards and should probably include audit and quality assurance processes, as well as realistic stopping rules to address unexpected or aberrant outcomes.

Symposium: Selection of patients for proton therapy

SP-0009

Patient selection for proton therapy: a clinicians view

A. Mahajan¹

¹MD Anderson Cancer Center, Proton Therapy Center, Houston, USA

Proton therapy is a radiation modality that has become increasingly available world wide over the past decade. It is an attractive radiotherapy intervention because of the charged particles dose deposition profile of characterized by the Bragg peak. By using proton therapy strategically, there is the possibility to deliver effective radiation dose to the target while reducing radiation to the surrounding non-target structures. The goals of any radiotherapy approach is to improve tumor control and/or reduce side effects and proton therapy offers an opportunity to achieve either one or both of these goals. Despite the promise of proton therapy, one must consider its associated risks and benefits, and as with any other radiation approach, to maximize the benefit to the patient. In general concepts that are useful in selecting and predicting a the benefit of proton therapy in individual patients include the following:

- 1) Proton therapy has the same risk of injury within the target area and high dose as other radiation therapies. For infiltrative tumors that require irradiation of a margin of normal tissue (example rhabdomyosarcoma) or those that have normal cells embedded within the tumor (example low grade glioma), the tissues receiving the high dose of radiotherapy will have similar risks of injury as non-proton approaches; therefore, one would not expect a lower risk of injury in the high dose area.
- 2) Since proton therapy is typically associated with a lower risk of late effects Patient who has a very low chance of surviving a long time due to the natural history of the disease, may not benefit from proton therapy, example widely metastatic cancer.
- 3) Patients, for example children, who can derive benefit from normal tissue radiation dose reduction are usually good candidates
- 4) Patients who require high doses of radiation to achieve tumor control, but would otherwise be limited due to normal tissue tolerance, for example patients with skull base chordoma or primary or secondary liver.
- 5) Tumor geometry and surrounding anatomy must be evaluated to estimate the potential benefit of proton therapy. For example, a 2 year old patient requiring flank radiation for Wilms tumor may have not benefit with proton therapy, whereas an 18 year old with a paravertebral Ewing's sarcoma may have significant advantage with proton therapy.
- 6) Patient set up, tissue uncertainties, external devices or implanted need to be evaluated to minimize the risk of uncertainties and disruption in the proton dosimetry.
- 7) Proton therapy may be a good option for re-irradiation in selected patients. In summary, proton therapy can be an excellent option to provide better local control and/or reduced toxicities in selected patients.

SP-0010

Selection of patients for proton therapy: a physicists view

M. Hoogeman¹, T. Arts¹, S. Van de Water¹, S. Van der Voort¹, Z. Perko², D. Lathouwers², S. Breedveld¹, B. Heijmen¹

¹Erasmus Medical Center Rotterdam, Erasmus MC Cancer Center, Rotterdam, The Netherlands

²Delft University of Technology, Radiation Science and Technology, Delft, The Netherlands

Intensity Modulated Proton therapy (IMPT) is a highly promising approach for radiation treatment of cancer patients due to its increased potential to reduce side effects and improve quality of life compared to contemporary radiation therapy techniques, such as IMRT. However, IMPT is associated with high costs and hence limited availability. Ideally, patient selection for IMPT should be based on the highest expected complication reduction compared to IMRT.

For a given patient, it is possible to predict the risk of side effects for proton and photon therapy by applying Normal Tissue Complication Probabilities (NTCP) models to optimized dose distributions. Only patients with clinically relevant reductions in NTCP exceeding minimum pre-defined thresholds will then qualify for proton therapy. While this approach should guarantee effective use of proton therapy, there are several concerns that will be discussed in this presentation:

1. The generation of a radiotherapy treatment plan is a complex procedure and its quality is highly dependent on the planner skills. To enable unbiased comparisons between IMPT and IMRT for each patient, automation of the treatment planning process is imperative.
2. IMPT is highly susceptible to inaccuracies in patient setup, anatomic changes, and to uncertainties in the calculation of the proton range. In IMRT, uncertainties in dose delivery are accounted for in the CTV-to-PTV margin. In IMPT, however, the PTV concept is not applicable. Alternatively, robust treatment planning can be used to take into account patient setup and range uncertainties. However, it is currently unknown which robustness settings need to be used to achieve an adequate target coverage for given population-based distributions of setup and range errors.
3. Image-guidance technology improves the accuracy of radiation therapy delivery, however its impact and current state-of-the-art may vary for proton and photon radiotherapy due to the physical differences between protons and photons and for historical reasons. The applied image-guidance technology will have an impact on the magnitude of NTCP reduction and hence on the selection of patients qualifying for proton therapy.

SP-0011

Future selection practice for proton therapy: selection of patients based on treatment planning comparison and NTCP-modelling

H. Langendijk¹

¹University Medical Center Groningen, Department of Radiation Oncology, Groningen, The Netherlands

The last decade, many new radiation delivery techniques have been clinically introduced without being subjected to randomized controlled trials. Many of these new techniques have been introduced in order to reduce the dose to the healthy tissues and subsequently to prevent radiation-induced side effects. Due to its superior beam properties, radiotherapy with protons compared to photons enables similar dose administration to the target volume with substantially lower dose to the normal tissue. In the Netherlands, we applied a 4-step model-based approach to select patients for proton therapy and to validate the benefit of protons compared to photons with regard to reducing the risk on radiation-induced side effects.

Step 1 consists of the development and validation of multivariable Normal Tissue Complication Probability (NTCP) models. NTCP models describe the relationship between radiation dose distribution parameters and the probability of a given side effect (NTCP-value). One of the output parameters of this step are the most relevant Dose Volume Histogram (DVH) parameters that can be used to optimize radiation treatment.

Step 2 includes in silico planning comparative studies. In this phase protons are compared with photons with regard to their ability to reduce the most relevant DVH-parameters resulting from step 1 (Δ Dose).

Step 3: Integration step 1 and 2. By integrating the results of the individual in silico planning comparison into the validated NTCP-models, the differences in dose can be translated into a difference in NTCP-value in each individual patient