

Soft Tissue Thickness Estimation for Head, Face, and Neck from CT Data for Product Design Purposes

Yang, W.; He, Renke; Chen, Tingshu; Goossens, R.H.M.; Huysmans, T.

DOI

[10.54941/ahfe1001983](https://doi.org/10.54941/ahfe1001983)

Publication date

2022

Document Version

Final published version

Published in

Ergonomics in Design: the magazine of human factors applications

Citation (APA)

Yang, W., He, R., Chen, T., Goossens, R. H. M., & Huysmans, T. (2022). Soft Tissue Thickness Estimation for Head, Face, and Neck from CT Data for Product Design Purposes. *Ergonomics in Design: the magazine of human factors applications*, 47, 558-565. <https://doi.org/10.54941/ahfe1001983>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Soft Tissue Thickness Estimation for Head, Face, and Neck from CT Data for Product Design Purposes

Wenxiu Yang¹, Renke He¹, Tingshu Chen¹, Haining Wang^{1*},
Richard Goossens², and Toon Huysmans²

¹School of Design, Hunan University, Changsha, 410000, China

²The Faculty of Industrial Design Engineering, Delft University of Technology,
Delft 2628CE, The Netherlands

ABSTRACT

Analysing the contact area between head-related products and the corresponding craniofacial profile is necessary to make such products more comfortable. The study of soft tissue biomechanical experiments provides a reliable perspective for understanding such contact areas and improving the comfort. In order to obtain more accurate and visualized craniofacial biological information that can be used to guide product design, CT data of the head, face, and neck of 50 Chinese aged 18-35 years were obtained in this paper. For each subject, an individual thickness map is calculated by segmentation of the soft tissue layer and wall-thickness calculation via Mimics. The individual maps superimposed on the outer surface area of the head, face, and neck are brought into correspondence using a non-rigid iterative closest point technique. From the correspondence an average head, face, and neck geometry and soft tissue thickness map was calculated. Statistics of the overall soft tissue thickness of the head, face, and neck is extracted, and an accurate soft tissue thickness map of the Chinese head, face, and neck is generated. This study not only lays the groundwork for future simulation experiments on head-related product design, but it also has significant implications for the fields of facial reconstruction in China.

Keywords: Product design, Comfort, Soft tissue thickness, CT scan

INTRODUCTION

“Comfort” is a key indicator in the field of product design, particularly in the design process of wearable products, to evaluate whether a product can achieve a good user experience while performing its basic functions. And “pressure” is one of the key factors that cause discomfort in wearable products. Researchers have shown that the constant application of force to a specific area can cause spots, rashes, skin irritations, or ulcers on the skin (Callaghan et al. 1998). Zhuang’s research also pointed out that if a device needs to be worn continuously for a long time, the pressure will increase with the wearing time (Dai et al. 2001), which leads to an increase in discomfort. This means that the degree of deformation of the soft tissue under force will affect the user’s assessment of the comfort.

However, the application of soft tissue deformation research in the field of ergonomics has not received extensive much attention. Most of the research on soft tissue is to solve medical problems, such as orthodontic surgery (Gisotti et al. 2022), tumor tissue (Hayes et al. 2004) and other biomechanical properties research. Luximon (Shah et al. 2021) has developed soft tissue deformation maps of pressure discomfort threshold (PDT) and pressure pain threshold (PPT) of the Chinese head and face in the field of ergonomics research. The degree of soft tissue deformation is proportional to the thickness of the corresponding soft tissue. It can be seen that in order to improve the pressure comfort of wearable products, anatomical and biomechanical studies on the soft tissues of the corresponding parts are critical. Because the focus of this research is on the head, face and neck, and the target product is the head-mounted product, the research will concentrate on soft tissue of the craniofacial region.

A large number of studies have measured the soft tissue thickness of the head and face among the existing facial soft tissue measurement studies. The facial soft tissue thickness data of Northern China (Shui et al. 2013), South-Korean (Cha et al. 2013), Brazilian (Tedeschi et al. 2009), White (Sahni et al. 2008) have been collected and analyzed. However, these studies only measure the thickness of soft tissue at a limited number of landmarks in the craniofacial region. There is no way to obtain accurate soft tissue thickness data of the head and face with a limited number of landmark measurements because there are countless feature points on the craniofacial part. Furthermore, the soft tissue thickness data obtained in the preceding studies are primarily presented in the form of data tables, which cannot directly display the thickness of the soft tissue of the head and face, nor can it be effectively applied in the design of head-mounted products.

Furthermore, in terms of measurement technology and application, traditional soft tissue thickness measurements typically use the needle depth probing method on cadavers (He et al. 2021). However, the measurement method and the cadaver's condition have a significant impact on the accuracy of the measurement results. This problem is effectively solved by the use of medical imaging techniques such as Computerized Tomography (CT).

Because of the scarcity of accurate and well-illustrated Chinese head and face soft tissue thickness structural data. CT data of the head, face, and neck of 50 Chinese aged 18–35 years were obtained using CT measurement technology in this paper. Later, the overall soft tissue thickness of the head, face, and neck is extracted, and an accurate soft tissue thickness map of the Chinese head, face, and neck is created. This study lays the groundwork for the head-related ergonomic product design.

MATERIAL AND METHODS

Materials

In this study, 50 subjects (24 males and 26 females) were selected for CT scans (the resolution of each CT data is $0.49 \times 0.49 \times 0.625$) at the Third Xiangya Hospital of Central South University. All of the study participants were Han Chinese from China's Hunan Province. Between the ages of 18




Figure 1: Soft tissue structure in Mimics (Green part represents soft tissue, blue part represents bone).




Figure 2: Model with brain (left) and without brain (right).

and 35. Subjects with a history of facial fractures, swelling, deformity, distortion, and pathological asymmetry, or any other pathology that might affect normal craniofacial features prior to measurement were barred from participating in the study. Before participation, all samples were given informed consent.

Methodology

The model processing procedure is as follows. The acquired CT data was first processed using Mimics 20.0 software in this study. The scan data is imported into the Mimics 20.0 software, the skull and soft tissue were separated using the system default threshold. Soft tissue Hounsfield units range from -700 to 225 . Figure 1 shows the soft tissue in green and the skull in blue. Then, since the focus of this study is on soft tissue and the skull, and a large




Figure 3: Soft tissue model with soft tissue thickness color coded for individual subject.



Figure 4: Skn part, top (left) and bottom (right) view.

number of intracranial structures will affect the speed of later model processing, most of the intracranial structures and band need to be removed at this stage (Figure 2).

After completing the model processing in Mimics 20.0 software, the model needs to be imported into 3-matic Research 12.0 for the calculation of the overall thickness of soft tissue (Figure 3). After that, the model and soft tissue thickness data were exported separately. Furthermore, all CT head models were extracted skin parts for importing in Paraview (Figure 4). R3DS Wrap (Dyke et al. 2020) was used to establish a correspondence between a template head model and the skin surface of the individual subjects (Figure 5).




Figure 5: CT headform (skin) and template headform after alignment (left), CT head model is being wrapped by the template head model (middle), Computed model (right).




Figure 6: Soft tissue thickness map (unisex).

The resulting homological head representations for each subject were equipped with individual soft tissue values at all vertices, derived from the soft



a. Female**b. Male**

Figure 7: Soft tissue thickness map for male and female.

tissue thickness that was calculated in 3-matic. The resulting homological head models could then be averaged in geometry and soft-tissue thickness. In addition, the head model was clipped to that region for which thickness data was available for a minimum of 40 subjects (Figures 6 and 7).

RESULT**Soft Tissue Thickness Map With High Accuracy**

Overall, soft tissue thickness is significantly thinner in regions with skull or teeth support compared to another craniofacial region. Furthermore, compared to the head and face, the neck, particularly the back of the neck, has the thickest soft tissue. In contrast, the soft tissue thickness of the scalp is the thinnest (Figures 6 and 7). Moreover, the soft tissue thickness for male is thicker than female ($p < 0.05$) (Table 1).

Table 1. Soft tissue thickness for 34 landmarks.

Landmark Number	Female Soft tissue thickness (mm) Mean	Male Soft tissue thickness (mm) Mean	Landmark Number	Female Soft tissue thickness (mm)	Male Soft tissue thickness (mm)
1	4.3	4.98	18	7.49	7.64
2	4.04	4.53	19	11.22	10.54
3	5.11	5.55	20	17.66	18.89
4	5.93	6.17	21	20.69	22.72
5	5.12	5.94	22	28.13	32.42
6	2.43	2.67	23	21.99	22.76
7	7.06	8.22	24	6.75	7.92
8	9.61	11.28	25	17.94	19.69
9	8.98	9.93	26	6.38	7.07
10	9.86	9.93	27	21.68	29.22
11	9.19	7.75	28	7.87	8.76
12	8.78	10.63	29	3.69	3.83
13	3.84	4.34	30	9.67	11.27
14	3.52	4.05	31	7.89	7.73
15	5.26	5.51	32	7.01	7.43
16	4.6	5.47	33	7.8	8.92
17	10.98	13.19	34	6.95	8.76

Application of Soft Tissue Research in Head-Related Product Design

The craniofacial soft tissue data obtained in this experiment can be used to create a model in the simulation experiment, which can then be used to run the simulation experiment. The traditional large-scale recruitment of users to evaluate the fit and comfort of the head-related product is relatively expensive, while the user test results of a small sample are not representative. The simulation experiment such as finite element force simulation experiment and fluid mechanics simulation experiment has many advantages. The simulation experiment can not only control the experiment cost, but it can also obtain specific information about each point of product, which has a wide range of applications in head-related product design.

CONCLUSION

3D anthropometric data are commonly used in ergonomics for comfort studies. However, this method is difficult to obtain specific and objective information of the contact part between the product and the human body. The biological characteristics of soft tissue can be further analyzed through the study of soft tissue anatomy. For example, the data collection in our paper, not only looking at averages but variations in soft tissue distribution by doing a Principal components analysis on the soft tissue thickness maps over the sample, which is beneficial to improving the design of head-related products.

ACKNOWLEDGMENT

The authors would like to acknowledge the support of School of design Hunan University, the Third Xiangya Hospital of Central South University and the Faculty of Industrial Design Engineering, Delft University of Technology.

REFERENCES

- Callaghan, S., & Trapp, M. (1998). Evaluating two dressings for the prevention of nasal bridge pressure sores. *Professional nurse (London, England)*, 13(6), 361–364.
- Cha, K. S. (2013). Soft-tissue thickness of South Korean adults with normal facial profiles. *The Korean journal of orthodontics*, 43(4), 178–185.
- Dai, J., Yang, J. J., & Zhuang, Z. (2011). Sensitivity analysis of important parameters affecting contact pressure between a respirator and a headform. *International Journal of Industrial Ergonomics*, 41(3), 268–279.
- Dyke, R. M., Lai, Y. K., Rosin, P. L., Zappalà, S., Dykes, S., Guo, D., ... & Yang, J. (2020). Shrec'20: Shape correspondence with non-isometric deformations. *Computers & Graphics*, 92, 28–43.
- Gisotti, M., Lops, D., & Valente, N. A. (2022). Esthetic management of space and soft tissue deficiencies in implant therapy: combined orthodontic-mucogingival approach. *The international journal of esthetic dentistry*, 17(1), 42–56.
- Hayes, A. J., Alexander, N., Clark, M. A., & Thomas, J. M. (2004). Elastofibroma: a rare soft tissue tumour with a pathognomonic anatomical location and clinical symptom. *European Journal of Surgical Oncology (EJSO)*, 30(4), 450–453.
- He, R., Yang, W., Wu, D., & Wang, H. (2021). Age- and Sex-Related Measurements of Total Craniofacial Soft Tissue Thickness and Fat in a Central Chinese Population. *The Journal of craniofacial surgery*, 32(8), 2626–2630.
- Sahni, D., Singh, G., Jit, I., & Singh, P. (2008). Facial soft tissue thickness in northwest Indian adults. *Forensic science international*, 176(2-3), 137–146.
- Shah, P., & Luximon, Y. (2021). Assessment of pressure sensitivity in the head region for Chinese adults. *Applied ergonomics*, 97, 103548.
- Shui W, Zhou M, Ji Y, et al. (2013). Measurement of facial soft tissue thickness and application to facial reconstruction. *Acta Anthropologica Sinica*. 2013;32: 345–353.
- Tedeschi-Oliveira, S. V., Melani, R. F. H., de Almeida, N. H., & de Paiva, L. A. S. (2009). Facial soft tissue thickness of Brazilian adults. *Forensic science international*, 193(1-3), 127–e1.