



Delft University of Technology

Integer packing sets form a well-quasi-ordering

Del Pia, Alberto; Gijswijt, Dion; Linderoth, Jeff; Zhu, Haoran

DOI

[10.1016/j.orl.2021.01.013](https://doi.org/10.1016/j.orl.2021.01.013)

Publication date

2021

Document Version

Accepted author manuscript

Published in

Operations Research Letters

Citation (APA)

Del Pia, A., Gijswijt, D., Linderoth, J., & Zhu, H. (2021). Integer packing sets form a well-quasi-ordering. *Operations Research Letters*, 49(2), 226-230. <https://doi.org/10.1016/j.orl.2021.01.013>

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.

Integer packing sets form a well-quasi-ordering

Alberto Del Pia ^{*} Dion Gijswijt [†] Jeff Linderoth [‡] Haoran Zhu [§]

January 17, 2021

Abstract

An integer packing set is a set of non-negative integer vectors with the property that, if a vector x is in the set, then every non-negative integer vector y with $y \leq x$ is in the set as well. The main result of this paper is that integer packing sets, ordered by inclusion, form a well-quasi-ordering. This result allows us to answer a recently posed question: the k -aggregation closure of any packing polyhedron is again a packing polyhedron.

Key words: Well-quasi-ordering; k -aggregation closure; polyhedrality; packing polyhedra.

1 Introduction

In order theory, a *quasi-order* is a binary relation \preceq over a set X that is *reflexive*: $\forall a \in X, a \preceq a$, and *transitive*: $\forall a, b, c \in X, a \preceq b$ and $b \preceq c$ imply $a \preceq c$. A quasi-order \preceq is a *well-quasi-order* (*wqo*) if for any infinite sequence x_1, x_2, \dots of elements from X there are indices $i < j$ such that $x_i \preceq x_j$.

A classic example of a quasi-order over the set of graphs is given by the graph minor relation. The Robertson-Seymour Theorem (also known as the graph minor theorem) essentially states that the set of finite graphs is well-quasi-ordered by the graph minor relation. This fundamental result is the culmination of twenty papers written as part of the Graph Minors Project [14]. Interested readers may find more examples and characterizations in the comprehensive survey paper by Kruskal [11]. The main result of this paper is that a quasi-order arising from Integer Optimization is a well-quasi-order.

Let $\mathbb{N} = \{0, 1, 2, \dots\}$ denote the set of nonnegative integers and let $[n] = \{1, 2, \dots, n\}$ for any $n \in \mathbb{N}$. We define an *integer packing set* in \mathbb{R}^n as a subset Q of \mathbb{N}^n with the property that: if $x \in Q$, $y \in \mathbb{N}^n$ and $y \leq x$, then $y \in Q$. Note that the relation \subseteq is a quasi-order over the set of integer packing sets. We are now ready to state our main result.

Theorem 1. *The set of integer packing sets in \mathbb{R}^n is well-quasi-ordered by the relation \subseteq .*

^{*}Department of Industrial and Systems Engineering & Wisconsin Institute for Discovery, University of Wisconsin-Madison. E-mail: delpia@wisc.edu.

[†]Delft Institute of Applied Mathematics, Delft University of Technology. E-mail: d.c.gijswijt@tudelft.nl.

[‡]Department of Industrial and Systems Engineering, University of Wisconsin-Madison. E-mail: linderoth@wisc.edu.

[§]Corresponding author. Department of Industrial and Systems Engineering, University of Wisconsin-Madison. E-mail: hzhu94@wisc.edu.

Integer packing sets appear naturally in Integer Optimization. A *packing polyhedron* is a set of the form $P = \{x \in \mathbb{R}^n \mid Ax \leq b, x \geq 0\}$ where the data $A \in \mathbb{R}^{n \times m}, b \in \mathbb{R}^m$ is non-negative. Clearly, for any packing polyhedron P , the set $P \cap \mathbb{Z}^n$, is an integer packing set. However, note that not all integer packing sets are of this form. This connection between packing polyhedra and integer packing sets allows us to employ Theorem 1 to answer a recently posed open question in Integer Optimization.

In [3], the authors introduce the concept of k -aggregation closure for packing and covering polyhedra. Given a packing polyhedron $P = \{x \in \mathbb{R}^n \mid Ax \leq b, x \geq 0\}$, and a positive integer k , the k -aggregation closure of P is defined by

$$\mathcal{A}_k(P) := \bigcap_{\lambda^1, \dots, \lambda^k \in \mathbb{R}_+^m} \text{conv}(\{x \in \mathbb{N}^n \mid (\lambda^j)^\top Ax \leq (\lambda^j)^\top b, \forall j \in [k]\}).$$

The set $\mathcal{A}_k(P)$ is defined as the intersection of an infinite number of sets, each of which is the convex hull of an integer packing set. A natural question, posed in [3], is whether the set $\mathcal{A}_k(P)$ is polyhedral. The authors provide a partial answer to this question by showing that $\mathcal{A}_k(P)$ is a polyhedron, provided that every entry of A is positive. As a consequence of Theorem 1, we give a complete answer to the posed question.

Theorem 2. *For any packing polyhedron P and any $k \geq 1$, the set $\mathcal{A}_k(P)$ is a packing polyhedron.*

The generality of our proof techniques allows us to provide a generalization of Theorem 2 to the setting where the given set is a downset of \mathbb{R}_+^n instead of a polyhedron. We recall that a *downset* of \mathbb{R}_+^n is a subset D of \mathbb{R}_+^n with the property that, if $x \in D, y \in \mathbb{R}_+^n$ and $y \leq x$, then $y \in D$. Clearly, a packing polyhedron in \mathbb{R}^n is a downset of \mathbb{R}_+^n , but not all downsets are polyhedral. Our generalization relies on a natural extension of the definition of k -aggregation closure to downsets of \mathbb{R}_+^n . For any downset D of \mathbb{R}_+^n , we denote by

$$\Lambda(D) := \{f \in \mathbb{R}^n \mid \sup\{f^\top x \mid x \in D\} < \infty\}.$$

In particular, note that $f^\top x \leq \beta$ is valid for D if and only if $f \in \Lambda(D)$ and $\beta \geq \sup\{f^\top x \mid x \in D\}$. Then, the k -aggregation closure of D is defined by

$$\tilde{\mathcal{A}}_k(D) := \bigcap_{f^1, \dots, f^k \in \Lambda(D)} \text{conv}(\{x \in \mathbb{N}^n \mid (f^j)^\top x \leq \sup\{(f^j)^\top d \mid d \in D\}, \forall j \in [k]\}).$$

The next observation shows that $\tilde{\mathcal{A}}_k$ is indeed a generalization of \mathcal{A}_k .

Observation 1. *For any packing polyhedron P and any $k \geq 1$, we have $\tilde{\mathcal{A}}_k(P) = \mathcal{A}_k(P)$.*

Proof. Let $P = \{x \in \mathbb{R}_+^n \mid Ax \leq b, x \geq 0\}$ be a packing polyhedron in \mathbb{R}^n . It is simple to show that $\tilde{\mathcal{A}}_k(P) \subseteq \mathcal{A}_k(P)$. To see this, consider an inequality $\lambda^\top Ax \leq \lambda^\top b$, for $\lambda \in \mathbb{R}_+^m$, in the definition of $\mathcal{A}_k(P)$. Then $\lambda^\top Ax \leq \lambda^\top b$ is valid for P . Thus $\lambda^\top A \in \Lambda(D)$, and $\sup\{\lambda^\top Ad \mid d \in P\} \leq \lambda^\top b$. Hence, the inequality $\lambda^\top Ax \leq \sup\{\lambda^\top Ad \mid d \in P\}$ in the definition of $\tilde{\mathcal{A}}_k(P)$ implies the original inequality $\lambda^\top Ax \leq \lambda^\top b$.

Next, we show $\tilde{\mathcal{A}}_k(P) \supseteq \mathcal{A}_k(P)$. Consider an inequality $f^\top x \leq \sup\{f^\top d \mid d \in P\}$, for $f \in \Lambda(P)$, in the definition of $\tilde{\mathcal{A}}_k(P)$. This inequality is valid for P . From Farkas' lemma we know that there exist some $\lambda \in \mathbb{R}_+^m$ and $\gamma \in \mathbb{R}_+^n$ such that $\lambda^\top A - \gamma^\top I = f^\top$ and $\lambda^\top b \leq \sup\{f^\top d \mid$

$d \in P\}$. Note that the inequality $\lambda^\top Ax \leq \lambda^\top b$ is valid for P . Furthermore, it dominates the inequality $f^\top x \leq \sup\{f^\top d \mid d \in P\}$ in the nonnegative orthant. This is because, whenever $x \geq 0$,

$$f^\top x = \lambda^\top Ax - \gamma^\top Ix \leq \lambda^\top Ax \leq \lambda^\top b \leq \sup\{f^\top d \mid d \in P\}.$$

We have shown $\tilde{\mathcal{A}}_k(P) \supseteq \mathcal{A}_k(P)$, which completes the proof of the observation. \square

We now state our generalizations of Theorem 2 to downsets of \mathbb{R}_+^n .

Theorem 3. *For any downset D of \mathbb{R}_+^n and any $k \geq 1$, the set $\tilde{\mathcal{A}}_k(D)$ is a packing polyhedron.*

In the special case $k = 1$, the k -aggregation closure is also known as the *aggregation closure*. In the recent unpublished manuscript [13], the authors independently show that the aggregation closure of a packing or covering rational polyhedron P is polyhedral. The main differences with our Theorem 3 are the following: (i) The result in [13] holds for both packing and covering polyhedra, while our Theorem 2 only deals with the packing case; (ii) The result in [13] requires the given set to be a polyhedron, while in our case the given set can be a general downset of \mathbb{R}_+^n ; (iii) The proof in [13] is direct, while our Theorem 2 is a consequence of Theorem 1; (iv) In [13] the authors only discuss in detail the aggregation closure, and claim that an analogous proof can be obtained for the k -aggregation closure, while in this paper we directly consider the k -aggregation closure.

Our techniques also allow us to obtain the following result.

Theorem 4. *For any closed convex downset D of \mathbb{R}_+^n , the set $\text{conv}(D \cap \mathbb{Z}^n)$ is a packing polyhedron.*

Our work sheds light onto the connection between Order Theory and polyhedrality of closures in Integer Optimization. Only few papers so far have explored this connection. In [2], Averkov exploits the Gordan-Dickson lemma to show the polyhedrality of the closure of a rational polyhedron obtained via disjunctive cuts from a family of lattice-free rational polyhedra with bounded max-facet-width. In the paper [7], Dash et al. consider fairly well-ordered qosets to extend the result of Averkov. In particular, the authors prove the polyhedrality of the closure of a rational polyhedron with respect to any family of t -branch sets, where each set is the union of t polyhedral sets that have bounded max-facet-width. Other recent polyhedrality results in Integer Optimization include [1, 4, 9, 6, 5].

The organization of this paper is as follows: In Section 2 we present some preliminaries and notations from Order Theory that will be used in our proofs. In Section 3 we show Theorem 1, while in Section 4 we provide a proof of Theorem 2. In Section 5 we turn our attention to non-polyhedral sets and prove Theorem 3 and Theorem 4.

2 Preliminaries in Order Theory

Recall that a *quasi-order* is a binary relation \preceq over a set X that is reflexive and transitive. If $a \preceq b$, we also write $b \succeq a$. If $a \preceq b$ or $b \preceq a$, the elements a and b are said to be *comparable*. If both $a \preceq b$ and $b \preceq a$, then we write $a \sim b$ (which is an equivalence relation). A sequence x_1, x_2, \dots of elements from X is said to be *increasing* if $x_1 \preceq x_2 \preceq \dots$ and *decreasing* if $x_1 \succeq x_2 \succeq \dots$.

Most quasi-orders in this paper will in fact be *partial orders*, that is, they are *antisymmetric*: $a \preceq b$ and $b \preceq a$ imply $a = b$. In particular, we will consider the subset relation on \mathbb{R}^n (and

the induced partial order on integer packing sets), and the partial order on (\mathbb{N}^n, \leq) given by the component-wise comparison: $x \leq y$ if and only if $x_i \leq y_i$ for all $i \in [n]$.

A quasi-order (X, \preceq) is a *well-quasi-order* (wqo) if for any infinite sequence of elements x_1, x_2, \dots from X there are indices $i < j$ such that $x_i \preceq x_j$. A quasi-order (X, \preceq) is said to have the *finite basis property* if for all $X' \subseteq X$, there exists a finite subset $B \subseteq X'$ such that for every $x \in X'$ there is a $b \in B$ such that $b \preceq x$. The next result provides us with characterizations of well-quasi-orders.

Lemma 1 ([10, Theorem 2.1]). *Let (X, \preceq) be a quasi-order. The following statements are equivalent:*

- (i) (X, \preceq) is a wqo;
- (ii) (X, \preceq) has the finite basis property;
- (iii) every infinite sequence of elements from X has an infinite increasing subsequence.

Given two quasi-orders (X_1, \preceq_1) and (X_2, \preceq_2) , the *product quasi-order* is $(X_1 \times X_2, \preceq)$ where $(x_1, x_2) \preceq (y_1, y_2)$ if and only if $x_1 \preceq_1 y_1$ and $x_2 \preceq_2 y_2$.

Lemma 2. *Let (X_1, \preceq_1) and (X_2, \preceq_2) be wqo's. Then the product quasi-order is a wqo.*

The proof of this well-known fact follows easily from the equivalence of (i) and (iii) in Lemma 1: given an infinite sequence of elements in $X_1 \times X_2$, we can find an infinite subsequence for which the components in X_1 form an increasing sequence, and then a further subsequence in which also the components in X_2 form an increasing sequence. The resulting subsequence is an increasing subsequence in $X_1 \times X_2$.

Since (\mathbb{N}, \leq) is a wqo, the lemma implies that for any positive n the set \mathbb{N}^n is a wqo under the usual component-wise comparison.

Lemma 3 (Gordan-Dickson, [8]). *The poset (\mathbb{N}^n, \leq) is a wqo.*

Given a quasi order (X, \preceq) we denote by X^* the set of all finite sequences of elements from X . We define a quasi order \preceq^* on X^* by setting $(x_1, \dots, x_n) \preceq^* (y_1, \dots, y_m)$ if and only if there is a strictly increasing function $f : [n] \rightarrow [m]$ such that $x_i \preceq y_{f(i)}$ for all $i \in [n]$ (in particular, we require $n \leq m$). In this paper we will need the following generalization of the Gordon-Dickson lemma.

Lemma 4 (Higman's lemma, [10]). *Let (X, \preceq) be a wqo. Then (X^*, \preceq^*) is a wqo as well.*

3 Integer packing sets are well-quasi-ordered

In this section we prove our main result that integer packing sets in \mathbb{R}^n form a wqo under inclusion. The proof is based on the following lemma.

Lemma 5. *Let (X, \preceq) be a wqo. Define X^{**} to be the set of decreasing sequences in X :*

$$X^{**} = \{(x_0, x_1, \dots) \in X^{\mathbb{N}} : x_0 \succeq x_1 \succeq \dots\}.$$

*For $x, y \in X^{**}$ set $x \preceq^{**} y$ if $x_i \preceq y_i$ for all $i \in \mathbb{N}$. Then (X^{**}, \preceq^{**}) is a wqo.*

Proof. We start with the following claim.

Claim 1. *Let $x \in X^{**}$. There is a $k \in \mathbb{N}$ such that $x_k \sim x_\ell$ for all $\ell \geq k$.*

Proof of claim. Since X is a wqo, it follows by Lemma 1 that there is a finite subset $I \subseteq \mathbb{N}$ of indices such that for any $\ell \in \mathbb{N}$ there is an $i \in I$ with $x_\ell \succeq x_i$. Let k be the largest index in I . Consider any $\ell \geq k$. Since x_0, x_1, \dots is decreasing, we have $x_\ell \preceq x_k$, but also $x_\ell \succeq x_i \succeq x_k$ for some $i \in I$. Hence, $x_\ell \sim x_k$. \diamond

We call the smallest k as in the claim the *tail* of x . By Higman's lemma, it follows that the product quasi-order \preceq' on $X^* \times X$ is a wqo. Let $\phi : X^{**} \rightarrow X^* \times X$ be defined by $\phi(x) = ((x_0, \dots, x_{k-1}), x_k)$, where k is the tail of x . Let $x, y \in X^{**}$ and suppose that $\phi(x) \preceq' \phi(y)$. To complete the proof, it suffices to show that $x \preceq^{**} y$.

Let k and ℓ be the tails of x and y , respectively. Since $\phi(x) \preceq' \phi(y)$ we have a strictly increasing function $f : \{0, \dots, k-1\} \rightarrow \{0, \dots, \ell-1\}$ such that $x_i \preceq y_{f(i)}$ for all $i \in \{0, \dots, k-1\}$.

Since $y_0 \succeq y_1 \succeq \dots$ and $f(i) \geq i$ for all $i \in \{0, \dots, k-1\}$, we have $x_i \preceq y_{f(i)} \preceq y_i$ for all $i \in \{0, \dots, k-1\}$. Since $x_k \preceq y_\ell$ and $y_\ell \leq y_j$ for all $j \in \mathbb{N}$ (as ℓ is the tail of y), it follows that for all $i \geq k$ we have $x_i \leq x_k \leq y_\ell \leq y_i$. We conclude that $x_i \leq y_i$ for all $i \in \mathbb{N}$ and therefore that $x \preceq^{**} y$. \square

We will now prove Theorem 1: the set of integer packing sets in \mathbb{R}^n is a wqo under inclusion.

Proof of Theorem 1. The proof is by induction on n . The case $n = 1$ follows directly from the fact that (\mathbb{N}, \leq) is a wqo. For the induction step, we associate to any integer packing set $S \subseteq \mathbb{R}^{n+1}$ a sequence (S_0, S_1, \dots) of 'slices' by setting

$$S_i = \{(x_1, \dots, x_n) \in \mathbb{N}^n : (x_1, \dots, x_n, i) \in S\}.$$

As S is an integer packing set, it follows that the S_i are integer packing sets in \mathbb{R}^n and that $S_0 \supseteq S_1 \supseteq \dots$. For two packing sets S, T in \mathbb{R}^{n+1} we have $S \subseteq T$ if and only if for the corresponding slices we have $S_i \subseteq T_i$ for all $i \in \mathbb{N}$. Hence, the well-quasi-ordering of integer packing sets in \mathbb{R}^{n+1} follows from that of integer packing sets in \mathbb{R}^n by Lemma 5. \square

As a consequence to Theorem 1 we obtain the following structural result about integer packing sets. An *n-dimensional block* is a set of the form $X_1 \times \dots \times X_n$, where each X_i is equal to \mathbb{N} or to $[m]$ for some $m \in \mathbb{N}$.

Corollary 1. *Let Q be an integer packing set in \mathbb{R}^n . Then Q is the union of finitely many n-dimensional blocks.*

Proof. The proof is by induction on n . If $n = 1$, then any integer packing set in \mathbb{R}^n is an n -dimensional block. Now suppose that the statement holds for a given n and consider an integer packing set Q in \mathbb{R}^{n+1} . Define the n -dimensional slices $Q_i = \{(x_1, \dots, x_n) : (x_1, \dots, x_n, i) \in Q\}$ for all $i \in \mathbb{N}$. Then Q_0, Q_1, \dots is a decreasing sequence of integer packing sets in \mathbb{R}^n . Hence, by Theorem 1, there is a $k \in \mathbb{N}$ such that $Q_k = Q_\ell$ for all $\ell \geq k$. By assumption, each set Q_i is a union of finitely many n -dimensional blocks. Hence $Q_i \times \{0, 1, \dots, i\}$ is a union of finitely many $n+1$ -dimensional block for any $i = 0, 1, \dots, k-1$, and also $Q_k \times \mathbb{N}$ is a union of finitely many $n+1$ -dimensional blocks. Since

$$Q = (Q_k \times \mathbb{N}) \cup \bigcup_{i=0}^{k-1} Q_i \times \{0, 1, \dots, i\},$$

the result follows. \square

4 Polyhedrality of the k -aggregation closure

In this section we prove that the k -aggregation closure of a packing polyhedron is itself a packing polyhedron (Theorem 2). We will use some standard notation from polyhedral theory. In particular, given $A \subseteq \mathbb{R}^n$, we denote by $\text{conv}(A)$ the convex hull of A , and given a polyhedron $P \subseteq \mathbb{R}^n$, we denote by $P_I = \text{conv}(P \cap \mathbb{Z}^n)$ the integer hull of P . Given $a \in \mathbb{R}^n$, we define $a_+ \in \mathbb{R}^n$ by $(a_+)_i := \max\{0, a_i\}$ for all $i \in [n]$.

Lemma 6. *Let D be a downset of \mathbb{R}_+^n and let $a^\top x \leq \beta$ be a valid inequality for D . Then $a_+^\top x \leq \beta$ is valid for D .*

Proof. Let $x \in D$ and let $x' \in \mathbb{R}^n$ be defined by $x'_i = x_i$ if $a_i \geq 0$ and $x'_i = 0$ if $a_i < 0$. Since D is a downset, we have $x' \in D$. Hence, $a_+^\top x = a^\top x' \leq \beta$. \square

Lemma 7. *A polyhedron is a downset of \mathbb{R}_+^n if and only if it is a packing polyhedron.*

Proof. It is simple to see that every packing polyhedron is a downset of \mathbb{R}_+^n . For the converse implication, let P be a polyhedron that is a downset of \mathbb{R}_+^n . Then P can be written in the form

$$P = \{x \in \mathbb{R}^n \mid x \geq 0, \quad (a^i)^\top x \leq b_i, \quad i \in [m]\}.$$

Consider any inequality $(a^i)^\top x \leq b_i$. Since P is a downset, it follows from Lemma 6 that $(a_+^i)^\top x \leq b_i$ is valid for P . Moreover, the inequality $(a^i)^\top x \leq b_i$ is implied by the inequalities $(a_+^i)^\top x \leq b_i$ and $x \geq 0$.

It follows that

$$P = \{x \in \mathbb{R}^n \mid x \geq 0, \quad (a_+^i)^\top x \leq b_i, \quad i \in [m]\}.$$

Since $0 \in P$, it follows that $b_i \geq 0$ for every $i \in [m]$. We conclude that P is a packing polyhedron. \square

Lemma 8. *Let P be a packing polyhedron. Then the integer hull P_I is also a packing polyhedron.*

Note that in Lemma 8 we do not require the polyhedron P to be rational.

Proof. We can write $P = \{x \in \mathbb{R}^n \mid x \geq 0, (a^i)^\top x \leq b_i, i \in [m]\}$ where the a^i and b_i are nonnegative. Consider any of the inequalities $(a^i)^\top x \leq b_i$. We claim that, for all $i \in [m]$, there exist a nonnegative rational vector $c^i \leq a^i$ and a rational number $d_i \geq b_i$ such that

$$\{x \in \mathbb{N}^n \mid (a^i)^\top x \leq b_i\} = \{x \in \mathbb{N}^n \mid (c^i)^\top x \leq d_i\}. \quad (1)$$

To show the claim, we observe that since a^i is nonnegative, the set $\{(a^i)^\top x \mid x \in \mathbb{N}^n\} \cap (b_i, b_i + 1)$ is finite. Hence, there is a number $b' > b_i$ such that for all $x \in \mathbb{N}^n$ we have either $(a^i)^\top x \leq b_i$ or $(a^i)^\top x \geq b'$.

Let $\epsilon \in (0, 1)$ be such that $(1 + \epsilon)b_i < b'(1 - \epsilon)$. We can choose c^i to be a rational vector with $(1 - \epsilon)a^i \leq c^i \leq a^i$, and d_i to be a rational number with $b_i \leq d_i \leq (1 + \epsilon)b_i$. Now, the inclusion \subseteq in equality (1) is clear. The converse inclusion follows since for any $x \in \mathbb{N}^n$ we have

$$(c^i)^\top x \leq d_i \implies (1 - \epsilon)(a^i)^\top x \leq (1 + \epsilon)b_i \implies (a^i)^\top x < b' \implies (a^i)^\top x \leq b_i.$$

This concludes the proof of the claim.

Let $P' = \{x \in \mathbb{R}^n \mid x \geq 0, (c^i)^\top x \leq d_i, i \in [m]\}$. Then $P \cap \mathbb{N}^n = P' \cap \mathbb{N}^n$ and hence $P_I = P'_I$. By Meyer's theorem [12], the integer hull of a rational polyhedron is itself a polyhedron. Hence, $P_I = P'_I$ is a polyhedron.

It is clear that the polyhedron P_I is a downset of \mathbb{R}_+^n . Hence, by Lemma 7, P_I is a packing polyhedron. \square

Now we are ready to present the proof of Theorem 2.

Proof of Theorem 2. Let P be a packing polyhedron defined by $P = \{x \in \mathbb{R}^n \mid Ax \leq b, x \geq 0\}$, and let k be a positive integer. Denote by \mathcal{P} the collection of polyhedra of the form

$$\{x \in \mathbb{R}^n \mid x \geq 0, (\lambda^j)^\top Ax \leq (\lambda^j)^\top b, \quad \forall j \in [k]\},$$

for all possible $\lambda^1, \dots, \lambda^k \in \mathbb{R}_+^m$.

Since A is nonnegative, for every $Q \in \mathcal{P}$ the set $Q \cap \mathbb{N}^n$ is an integer packing set. By Theorem 1, the set of integer packing sets in \mathbb{R}^n is a wqo under inclusion. Hence, it follows from the finite basis property that there is a finite subset $\mathcal{P}' \subseteq \mathcal{P}$ such that for any $Q \in \mathcal{P}$ there is a $Q' \in \mathcal{P}'$ such that $Q' \cap \mathbb{N}^n \subseteq Q \cap \mathbb{N}^n$, and hence also that $Q'_I \subseteq Q_I$. We conclude that

$$\mathcal{A}_k(P) = \bigcap \{Q_I : Q \in \mathcal{P}\} = \bigcap \{Q_I : Q \in \mathcal{P}'\}.$$

Since by Lemma 8 the integer hull Q_I is a packing polyhedron for every $Q \in \mathcal{P}'$, and the intersection of finitely many packing polyhedra is again a packing polyhedron, it follows that $\mathcal{A}_k(P)$ is a packing polyhedron. \square

5 Generalization to non-polyhedral sets

In this section, we provide the proofs of our generalizations of Theorem 2 to non-polyhedral sets. In particular, we give the proofs of Theorem 3 and of Theorem 4.

Proof of Theorem 3. Define $\Lambda^+(D) := \Lambda(D) \cap \mathbb{R}_+^n$. We first show that in the definition of $\tilde{\mathcal{A}}_k(D)$ we can replace $\Lambda(D)$ with $\Lambda^+(D)$, i.e.,

$$\tilde{\mathcal{A}}_k(D) = \bigcap_{f^1, \dots, f^k \in \Lambda^+(D)} \text{conv}(\{x \in \mathbb{N}^n \mid (f^j)^\top x \leq \sup\{(f^j)^\top d \mid d \in D\}, \quad \forall j \in [k]\}).$$

The containment \subseteq is trivial, thus we only need to show the containment \supseteq . Let $f \in \Lambda(D)$, and consider the associated valid inequality for D given by $f^\top x \leq \sup\{f^\top d \mid d \in D\}$. Since D is a downset of \mathbb{R}_+^n , we know from Lemma 6 that $(f_+)^\top x \leq \sup\{f^\top d \mid d \in D\}$ is also valid for D , and dominates the original inequality in \mathbb{R}_+^n . In particular, this implies that $\sup\{(f_+)^\top d \mid d \in D\} \leq \sup\{f^\top d \mid d \in D\}$, hence $f_+ \in \Lambda^+(D)$ since the latter supremum is finite by assumption. Hence, we have shown that $(f_+)^\top x \leq \sup\{(f_+)^\top d \mid d \in D\}$ dominates the original inequality $f^\top x \leq \sup\{f^\top d \mid d \in D\}$ in \mathbb{R}_+^n . We have therefore proven the containment \supseteq .

Lastly, we follow almost the exact same argument as in the proof of Theorem 2, except now we consider the collection \mathcal{P} of polyhedra of the form

$$\{x \in \mathbb{R}^n \mid x \geq 0, (f^j)^\top x \leq \sup\{(f^j)^\top d \mid d \in D\}, \quad \forall j \in [k]\},$$

for all possible $f^1, \dots, f^k \in \Lambda^+(D)$. \square

We now turn our attention to Theorem 4.

Proof of Theorem 4. For any $f \in \Lambda(D)$ we denote $\beta_f := \max\{f^\top d \mid d \in D\}$ and define $\Lambda^+(D) := \Lambda(D) \cap \mathbb{R}_+^n$ as in the previous proof. We obtain

$$\begin{aligned} D &= \{x \in \mathbb{R}_+^n \mid f^\top x \leq \beta_f, \forall f \in \Lambda^+(D)\}, \\ \text{conv}(D \cap \mathbb{Z}^n) &= \text{conv}(\{x \in \mathbb{N}^n \mid f^\top x \leq \beta_f, \forall f \in \Lambda^+(D)\}). \end{aligned}$$

For any $f \in \Lambda^+(D)$ let $S_f := \{x \in \mathbb{N}^n \mid f^\top x \leq \beta_f\}$. Then S_f is an integer packing set in \mathbb{R}^n . By Theorem 1, the set of integer packing sets in \mathbb{R}^n is a wqo under inclusion. Hence, it follows from the finite basis property that there is a finite subset $B \subseteq \Lambda^+(D)$ such that for every $f \in \Lambda^+(D)$ there is a $f' \in B$ for which $S_{f'} \subseteq S_f$. It follows that

$$\text{conv}(D \cap \mathbb{Z}^n) = \text{conv}(\{x \in \mathbb{N}^n \mid f^\top x \leq \beta_f, \forall f \in B\}).$$

By Lemma 8, it follows that $\text{conv}(D \cap \mathbb{Z}^n)$ is a packing polyhedron. \square

Theorem 4 can be used to prove the following result about the natural extension of $\tilde{\mathcal{A}}_k$ to $k = \infty$ defined by

$$\tilde{\mathcal{A}}_\infty(D) := \text{conv}(\{x \in \mathbb{N}^n \mid f^\top x \leq \sup\{f^\top d \mid d \in D\}, \forall f \in \Lambda(D)\}).$$

Corollary 2. *For any downset D of \mathbb{R}_+^n , the set $\tilde{\mathcal{A}}_\infty(D)$ is a packing polyhedron.*

Proof. From Lemma 6, $\overline{\text{conv}}(D)$ is a closed convex downset, and we can write

$$\overline{\text{conv}}(D) = \{x \in \mathbb{R}_+^n \mid f^\top x \leq \sup\{f^\top d \mid d \in D\}, \forall f \in \Lambda(D)\}.$$

Hence $\tilde{\mathcal{A}}_\infty(D) = \text{conv}(\overline{\text{conv}}(D) \cap \mathbb{N}^n)$. The corollary then follows from Theorem 4. \square

Acknowledgments

The authors would like to thank the anonymous referee for suggesting that Theorem 4 was implied by our earlier results and for a careful reading of the manuscript.

References

- [1] K. Andersen, Q. Louveaux, and R. Weismantel. An analysis of mixed integer linear sets based on lattice point free convex sets. *Mathematics of Operations Research*, 35(1):233–256, 2010.
- [2] G. Averkov. On finitely generated closures in the theory of cutting planes. *Discrete Optimization*, 9(4):209–215, 2012.
- [3] M. Bodur, A. Del Pia, S. Dey, M. Molinaro, and S. Pokutta. Aggregation-based cutting-planes for packing and covering integer programs. *Mathematical Programming*, pages 1–29, 2017.

- [4] D. Dadush, S. Dey, and J. Vielma. On the Chvátal-Gomory closure of a compact convex set. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 130–142. Springer, 2011.
- [5] S. Dash, O. Günlük, and D. Lee. Generalized Chvátal-Gomory closures for integer programs with bounds on variables. *Mathematical Programming*, pages 1–33, 2020.
- [6] S. Dash, O. Günlük, and D. Morán. On the polyhedrality of cross and quadrilateral closures. *Mathematical Programming*, 160(1-2):245–270, 2016.
- [7] S. Dash, O. Günlük, and D. Morán. On the polyhedrality of closures of multibranch split sets and other polyhedra with bounded max-facet-width. *SIAM Journal on Optimization*, 27(3):1340–1361, 2017.
- [8] L. Dickson. Finiteness of the odd perfect and primitive abundant numbers with n distinct prime factors. *American Journal of Mathematics*, 35(4):413–422, 1913.
- [9] J. Dunkel and A. Schulz. The Gomory-Chvátal closure of a nonrational polytope is a rational polytope. *Mathematics of Operations Research*, 38(1):63–91, 2013.
- [10] G. Higman. Ordering by divisibility in abstract algebras. *Proceedings of the London Mathematical Society*, 3(1):326–336, 1952.
- [11] J. Kruskal. The theory of well-quasi-ordering: A frequently discovered concept. *Journal of Combinatorial Theory, Series A*, 13(3):297–305, 1972.
- [12] Robert R Meyer. On the existence of optimal solutions to integer and mixed-integer programming problems. *Mathematical Programming*, 7(1):223–235, 1974.
- [13] K. Pashkovich, L. Poirrier, and H. Pulyassary. The aggregation closure is polyhedral for packing and covering integer programs. *arXiv preprint arXiv:1910.03404v1*, 2019.
- [14] N. Robertson and P. Seymour. Graph minors. XX. Wagner’s conjecture. *Journal of Combinatorial Theory, Series B*, 92(2):325–357, 2004.