

Document Version

Final published version

Licence

Dutch Copyright Act (Article 25fa)

Citation (APA)

Kapanidis, A. N., Muras, L., Sreenivasa, K., Hazra, J. P., van Noort, J., Joo, C., & Deindl, S. (2026). From sequence to function: Bridging single-molecule kinetics and molecular diversity. *Science (New York, N.Y.)*, 391(6784), 457-465. <https://doi.org/10.1126/science.adv4503>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

In case the licence states "Dutch Copyright Act (Article 25fa)", this publication was made available Green Open Access via the TU Delft Institutional Repository pursuant to Dutch Copyright Act (Article 25fa, the Taverne amendment). This provision does not affect copyright ownership.
Unless copyright is transferred by contract or statute, it remains with the copyright holder.

Sharing and reuse

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

REVIEW SUMMARY

MOLECULAR BIOLOGY

From sequence to function: Bridging single-molecule kinetics and molecular diversity

A. N. Kapanidis*, L. Muras, K. Sreenivasa, J. P. Hazra, J. van Noort, C. Joo, S. Deindl*

Full article and list of author affiliations:
<https://doi.org/10.1126/science.adv4503>

BACKGROUND: Biomolecules derive their function from sequence. Small changes in DNA, RNA, or protein sequence can shift structure, mechanics, and reaction pathways. These effects shape folding, binding equilibria, and the mechanochemical cycles of molecular machines. Ensemble assays have mapped many sequence-function relationships, but population averaging obscures transient intermediates, rare states, alternative pathways, and static heterogeneity.

Single-molecule methods overcome this limitation. Techniques such as single-molecule fluorescence resonance energy transfer (smFRET), colocalization imaging, optical and magnetic tweezers, and nanopore-based force spectroscopy reveal real-time transitions between structural states that bulk measurements hide. Yet typical single-molecule experiments are slow and low throughput, typically probing one construct at a time. This bottleneck has limited our ability to explore large sequence spaces and to connect specific sequences to detailed kinetics.

ADVANCES: New multiplexed single-molecule strategies now address this gap. Recent work links the functional “phenotype” of an individual molecule to its precise sequence across thousands of variants in one experiment. Two general approaches have emerged: MUSCLE (multiplexed single-molecule characterization at the library scale) and SPARXS (single-molecule parallel analysis for rapid exploration of sequence space) couple TIRF (total internal reflection fluorescence)-based fluorescence trajectories with in situ Illumina sequencing, generating millions of spatially registered traces tied directly to a known variant. SPIN-seq (single-molecule phenotyping and in situ sequencing) uses single-molecule sequencing-by-hybridization on surface-immobilized DNA to identify each sequence variant after kinetic imaging on the same TIRF microscope. These methods now enable sequence-resolved measurements of conformational dynamics, kinetic landscapes, reaction pathways, and static heterogeneity at scale.

This capability has already revealed how sequence tunes Holliday junction isomerization, DNA hairpin breathing, and

Cas9-induced target unwinding and rewinding. It complements earlier single-molecule work showing that the stepping behavior of polymerases, helicases, polymerases, and ribosomes is dictated by local, sequence-dependent energy barriers. Multiplexing now generalizes such analyses to full libraries measured under identical conditions.


Technical challenges remain: Field-of-view limitations, photobleaching, and incomplete cluster formation reduce throughput relative to standard next-generation sequencing. Rare behaviors may be missed when sampling limits are reached, and nonequilibrium reactions are difficult to synchronize across large sample surfaces. Advances in imaging hardware, photostable labeling, and optically gated reaction initiation will be key for further development.

OUTLOOK: The scope of multiplexed single-molecule techniques is also expanding beyond nucleic acids. Display technologies or DNA-barcoded proteins could support high-throughput kinetic analysis of peptide and protein libraries, including protein-protein, protein-ligand, and aptamer-ligand interactions relevant to drug discovery. Integrations with super-resolution imaging, force spectroscopy, or DNA origami scaffolds may extend measurements to mesoscale (tens to hundreds of nanometers up to a few micrometers) structural motions and rotational degrees of freedom. The iterative interplay between large-scale single-molecule datasets and machine-learning models will enable deeper interpretation and accelerate the discovery of sequence-function relationships.

Ultimately, these developments reunite high-throughput sequencing with single-molecule biophysics. Tying molecular identity to dynamic behavior for millions of molecules provides a powerful route to map sequence-structure-function relationships with unprecedented resolution. This convergence promises a next generation of tools for mechanistic biology and molecular engineering. □

*Corresponding author. Email: kapanidis@physics.ox.ac.uk (A.N.K.); sebastian.deindl@icm.uu.se (S.D.)

Sequence-function relationships across complex kinetic landscapes. Conventional single-molecule assays reveal functional phenotypes such as Cas9 R-loop formation, rewinding, and conformational equilibria yet seldom show how these dynamic processes depend on sequence. Multiplexed approaches integrate pooled library assembly, kinetic imaging, in situ sequencing, and per-molecule registration to assign trajectories to specific variants, enabling direct analysis of how sequence modulates energetic landscapes, intermediate states, and pathway heterogeneity.



MOLECULAR BIOLOGY

From sequence to function: Bridging single-molecule kinetics and molecular diversity

A. N. Kapanidis^{1,2*}, L. Muras³, K. Sreenivasa⁴, J. P. Hazra^{1,2}, J. van Noort⁵, C. Joo^{4,6}, S. Deindl^{3*}

Biological function is fundamentally determined by nucleic acid and protein sequence. Beyond encoding genetic information, nucleic acids also display complex physicochemical parameters that shape structure, dynamics, and interactions. Understanding how sequence variation sculpts the energetic landscapes underlying these properties requires methods that capture both molecular diversity and dynamic behavior. Single-molecule techniques are ideally suited to this task, but conventional formats remain time and cost intensive. Recent breakthroughs have enabled highly multiplexed approaches for observing molecular dynamics across millions of individual molecules representing thousands of sequences or barcoded entities. Though still in development, these methods have begun to bridge sequence, structure, dynamics, and function at scale, opening new opportunities in drug discovery, molecular diagnostics, and functional genomics.

Biological function is fundamentally determined by the sequence of nucleic acids and proteins (Fig. 1). This principle is especially evident in the role of DNA and RNA in encoding genetic information, which in turn dictates the sequences and structural properties of proteins (Fig. 1A). Beyond serving as carriers of genetic instructions, nucleic acids also exhibit complex physicochemical properties that critically influence their structure, dynamics, and interactions (Fig. 1, B to D). Differences in the chemical properties of the DNA bases enable specific interactions with proteins, such as transcription factors, through contacts in both the major and minor grooves of the DNA (Fig. 1D).

Within double-stranded DNA, variations in hydrogen bonding between base pairs and differences in base-stacking interactions between the aromatic rings of adjacent bases make the mechanical properties of DNA sequence-dependent (Fig. 1, B and C). Such variations can have important biological consequences, facilitating melting bubbles that enable transcription and recombination, introducing bends and kinks, and giving rise to alternative structures such as G-quadruplexes. Furthermore, local variations in the mechanical properties of DNA suggest the existence of a mechanical code for positioning nucleosomes (1, 2), adding another layer of transcription regulation. RNAs, being single-stranded, allow for an even wider spectrum of sequence-dependent structures and dynamic conformational exchanges. As a result, DNA and RNA store and transmit information beyond the explicit genetic code. Achieving a comprehensive understanding of how sequence variations shape the energetic landscapes (Fig. 1E) that govern structural

dynamics and molecular interactions, and ultimately modulate biological function, often requires exploration of a broad sequence space.

Recent ensemble-based methods for high-throughput *in vitro* analysis have substantially advanced mechanistic molecular biology (2–19). Many of these approaches rely on DNA sequencing to distinguish between different library members. In these cases, DNA sequences can be read out directly or used as indirect proxies for RNA or protein sequences through transcription, translation, or DNA barcoding. For example, massively parallel reporter assays link DNA barcodes to regulatory elements to quantify their effects on gene expression (20, 21). Deep mutational scanning enables systematic mapping of the functional consequences of mutations across a protein or RNA sequence (22, 23). In another approach, DNA-barcoded protein-binding assays can reveal sequence determinants of transcription factor binding with high resolution (4, 24). More broadly, DNA barcoding extends these strategies to encompass diverse types of chemical variation (6, 7). Although these methods offer considerable advantages, their capacity to resolve dynamic processes is inherently constrained by ensemble averaging.

Single-molecule measurements and the power of multiplexing

Single-molecule techniques provide a powerful means to study these dynamic processes, as they circumvent ensemble averaging, which can obscure transient intermediates and alternative kinetic pathways (25–43). However, single-molecule methods are often limited in their ability to observe multiple distinct molecular species, typically requiring separate labor-intensive experiments for each species, one at a time. This constraint often restricts single-molecule investigations to a small subset of possible samples, making it impractical to study large libraries.

Recent studies have demonstrated the simultaneous analysis of mixtures containing two (44) or nine (45) distinct DNA sequences using fluorophore-labeled detection oligonucleotides. More recently, Squires *et al.* demonstrated an elegant solution-phase alternative in which nanostructured fluorescence resonance energy transfer (FRET) barcodes, held in an anti-Brownian electrokinetic trap, discriminate on the order of two dozen DNA species (46). Although the barcodes markedly expand the distinguishable palette, this approach does not provide a direct readout of each molecule's sequence and is presently limited to tens of variants. Collectively, these studies established a proof of concept for correlating the functional characteristics of an individual DNA molecule, referred to as its “single-molecule phenotype,” with its corresponding DNA sequence. This phenotype may encompass structural or dynamic features of molecular interactions, such as the kinetics of a biochemical reaction at the single-molecule level, the equilibrium dynamics of binding and unbinding events, or conformational changes in a protein-DNA complex. For surface-immobilized libraries, the workflow involves immobilization of a library of distinct DNA sequences on a solid support, a single-molecule fluorescence measurement that evaluates a single-molecule phenotype of each DNA molecule, and a step where the library complexity is interrogated by sequential transient DNA hybridization. By contrast, the solution-phase barcode strategy decouples the phenotype measurement from any surface attachment, relying instead on photophysical tag combinations for molecular identity. Both routes, however, still face scalability challenges when applied to truly large libraries.

In another study, single-molecule fluorescence imaging was combined with *in situ* single-molecule DNA sequencing to provide an endpoint measurement for identifying histone modifications and genomic positions of individual nucleosomes (47, 48). However, the protocol used did not capture single-molecule dynamics, limiting its application to mapping populations. These pioneering efforts relied on custom methodologies, and their progression is hampered by the absence of commercial platforms for *in situ* single-molecule sequencing.

Linking single-molecule kinetics to nucleic acid sequence

The development of DNA sequencing approaches has often been closely intertwined with advancements in single-molecule methods,

¹Department of Physics, University of Oxford, Oxford, UK. ²Kavli Institute for Nanoscience Discovery, University of Oxford, Oxford, UK. ³Department of Cell and Molecular Biology, Science for Life Laboratory, Uppsala University, Uppsala, Sweden. ⁴Department of BioNanoScience, Kavli Institute of Nanoscience, Delft University of Technology, Delft, Netherlands. ⁵Biological and Soft Matter Physics, Huygens-Kamerlingh Onnes Laboratory, Leiden University, Leiden, Netherlands. ⁶Department of Physics, Ewha Womans University, Seoul, Republic of Korea. *Corresponding author. Email: kapanidis@physics.ox.ac.uk (A.N.K.); sebastian.deindl@icm.uu.se (S.D.)




Fig. 1. Sequence variation modulates structure, dynamics, and interactions of biomolecules across multiple levels. (A) Schematic overview of the hierarchical information flow from DNA to folded protein according to the central dogma of molecular biology, illustrating how DNA sequence not only encodes amino acid identity but also shapes structural and dynamic properties at each stage of molecular processing. (B) The sequences of DNA and RNA encode their characteristic secondary and tertiary structures. Shown are canonical duplex forms such as B-DNA and A-DNA, as well as noncanonical motifs that include G-quadruplexes and structured aptamers. Common RNA secondary structures include stem-loop hairpins, whereas fully folded tRNAs and ribozymes exhibit complex tertiary architectures shaped by sequence context and base identity. [tRNA structure adapted from ToGo Picture Gallery, CC BY 4.0] (C) Sequence variation shapes energetic landscapes and transition kinetics of folding and unfolding pathways. Shown are examples of force-induced hairpin unfolding, DNA bubble formation, and conformational exchange within structured RNA or DNA segments, such as Holliday junctions (HJ). (D) DNA sequence controls interactions with proteins, including the sequence-dependent recognition by transcription factors and RNAP (top) as well as the sequence-dependent Cas9 binding and target DNA unwinding (bottom). Energy landscapes illustrate sequence-specific kinetic barriers and intermediates during enzyme function. Indicated are energy differences between bound and free states (including transition barriers) and the kinetic rates of target-site search by facilitated diffusion. [In vivo transcription factor binding energy landscape adapted from (125), CC BY 4.0; genome editing energy landscape adapted from (126), CC BY 4.0] (E) In complex multistep biological processes, different sequences give rise to distinct energy landscapes. Sequence variation reshapes both the geometry of the energy surface and the accessibility of kinetic pathways, favoring distinct major and minor pathways depending on barrier heights, cooperative transitions, and environmental constraints.

with each technology driving the other's growth and impact. For example, the pursuit of single-molecule DNA sequencing catalyzed early advances in single-molecule fluorescence spectroscopy in solution. Pioneering efforts in R. Keller's lab at Los Alamos involved attempts to sequence DNA by digesting immobilized, fluorescently labeled DNA fragments using an exonuclease and then detecting the cleaved fluorescent nucleotides as they passed through a laser-interrogated detection zone (49). Similarly, the original aim of the company Solexa in the 1990s was to achieve fluorescence-based single-molecule DNA sequencing. Although these initial efforts were hampered by sensitivity challenges, they ultimately led to the development of clonal-cluster DNA next-generation sequencing (NGS), later commercialized by Illumina.

Other efforts, motivated by the push for rapid and affordable genome sequencing during the Human Genome Project, were driven by single-molecule biophysics groups and led to several single-molecule technologies that made third-generation DNA sequencing (based on single-molecule detection) a reality. These efforts began with seminal proof-of-concept studies (Fig. 2) that demonstrated the feasibility of reading sequence information from single DNA molecules using either fluorescence- or force-based approaches (50–52), followed by the first demonstration of single-molecule DNA sequencing (53) (Fig. 2, A and B). This work culminated in technologies such as PacBio sequencing, which relies on the real-time incorporation of labeled nucleotides in zero-mode waveguides (54) (Fig. 2C), and nanopore sequencing, which reads DNA sequence information by detecting electrical signal changes as individual bases of nucleic acid strands pass through a protein nanopore (55) (Fig. 2D).

The interplay between single-molecule methods and DNA sequencing, however, extends beyond decoding the one-dimensional sequence information for nucleic acids. From their inception, single-molecule experiments have explored the mechanical properties of biological heteropolymers, particularly DNA and RNA, and how these properties influence function and biological roles. This interest stemmed both from the rich information content of nucleic acids and from practical considerations, such as the ease of preparation, labeling, modification, and surface immobilization for single-molecule studies. From the earliest single-molecule FRET measurements (56, 57), researchers explored sequence variation and modifications, initially to validate methodologies, then to test hypotheses, and ultimately to uncover unexpected behaviors using powerful single-molecule techniques. Sequence dependence played a central role in many studies on ribozymes and RNA folding kinetics (57, 58), DNA hybridization (59), and protein-nucleic acid interactions, such as exonucleases and nucleic acid polymerases.

A key example involves the bacterial RNA polymerase (RNAP), which catalyzes DNA-templated RNA synthesis. Early single-molecule optical trapping experiments measured the force and velocity of RNAP during transcription on DNA and revealed that the RNAP motion on the DNA during transcription elongation (and the associated transcription kinetics) was influenced by DNA sequence (60). Subsequent studies demonstrated frequent polymerase pausing, independent of RNAP backtracking (61). Early single-molecule work by Greenleaf and Block (52) began to establish a link between RNAP pausing and DNA sequence (Fig. 2A), anticipating later genome-wide approaches. However, the sequence dependence of this ubiquitous pausing remained unclear until single-molecule measurements were combined with NGS (62, 63), leading to the identification of a consensus DNA sequence associated with frequent elongation pauses and to the mechanistic dissection of these events. The power of single-molecule approaches in probing sequence-dependent functional effects was further exemplified by recent high-resolution nanopore tweezers studies that track RNAP motion along the DNA template with unprecedented temporal and spatial resolution, resolving translocation events smaller than a single base-pair step (<3.4 Å). These studies revealed a complex kinetic landscape in which RNAP transitions between multiple transcriptional states, with these transitions being highly DNA-sequence dependent




Fig. 2. Early single-molecule approaches linking DNA sequence to molecular function.

(A) Motion-based DNA sequencing using optical tweezers. Greenleaf and Block (52,127) tracked single RNAP molecules transcribing in dual optical traps. Limiting the concentration of one type of nucleotide at a time caused site-specific pauses, producing distinct patterns that reveal sequence. Aligned transcription traces under different conditions generate position histograms, from which base identity is inferred. [Adapted with permission from (52); adapted with permission from Springer Nature (127)] (B) Early sequencing-by-synthesis with asynchronous base incorporation. Harris *et al.* (53) hybridized poly(dA)-tailed templates to surface-tethered poly(dT) primers. Each cycle added a single fluorescently labeled nucleotide, followed by polymerase extension, imaging, dye cleavage, and washing. Fluorescence at fixed positions across cycles allowed direct single-molecule reads, laying the groundwork for today's multiplexed platforms. [Adapted with permission from (53)] (C) Real-time sequencing in zero-mode waveguides (ZMWs). Eid *et al.* (54) used immobilized DNA polymerase and fluorescently labeled deoxynucleotide triphosphates (dNTPs) confined in ZMWs. Fluorophores on terminal phosphates allowed real-time detection without inhibiting enzymatic activity. Early traces show color-coded incorporation events across a ~ 28 -base read, capturing nucleotide identity and polymerase kinetics. Commercial implementations monitor tens of thousands of ZMWs in parallel, but a single ZMW is depicted for clarity. [Adapted with permission from (54)] (D) Nanopore-based methods linking sequence with dynamics. (i) Clarke *et al.* (55) demonstrated sequencing through a modified α -hemolysin pore with an adapter, detecting individual nucleotides through ionic current signatures, an early foundation for modern nanopore sequencing. Modern nanopore sequencers use large arrays of pores and molecular barcoding to achieve massive parallelization; here, a single pore is shown to illustrate the underlying principle. (ii) Nova *et al.* (64) combined nanopore sensing with force application. Using engineered templates and transcription complexes, RNAP was held at the pore rim while voltage-induced translocation revealed subnucleotide steps and sequence-dependent pausing, capturing both dynamic and mechanical features of transcription. [(i) Adapted with permission from Springer Nature (55); (ii) reproduced from (64), CC BY-NC-ND 4.0]

and modulated by interactions between DNA and structural elements of RNAP (64) (Fig. 2D). Nanopore tweezers strategies have also been applied to helicases, providing detailed mechanistic insights (65–67), and helicases of this type also serve as processive motors in nanopore sequencing platforms to regulate translocation through the pore (68). Extending beyond RNAP and helicases, similar advances have been achieved for other linear molecular machines that operate on nucleic acids, such as DNA polymerase (69) and the ribosome (70). However, the requirement for linear motion along a template constrains the scope of these technologies, necessitating the development of additional, more general methods to study the broader landscape of protein-DNA interactions and biomolecular interactions in general.

Multiplexed single-molecule measurements

In addressing the need for additional methods, two distinct strategies for highly multiplexed single-molecule measurements (see Box 1) have recently emerged: the closely related MUSCLE (multiplexed single-molecule characterization at the library scale) (71) and SPARXS (single-molecule parallel analysis for rapid exploration of sequence space) (72) methods as well as SPIN-seq (single-molecule phenotyping and in situ sequencing) (73) (Fig. 3).

Integrating single-molecule fluorescence with Illumina sequencing

The MUSCLE and SPARXS methods integrate single-molecule fluorescence microscopy with Illumina NGS (74) to monitor complex dynamic behaviors at the single-molecule level while subsequently determining their sequence (71, 72). The workflow begins with the design of a molecular library that includes sequencing adapters for immobilization on the surface of an Illumina MiSeq flow cell, ensuring compatibility with both single-molecule imaging and sequencing (Fig. 3A). The MiSeq flow cells are first examined by single-molecule fluorescence imaging of surface-bound constructs using objective-type total internal reflection fluorescence (TIRF) microscopy. With highly automated TIRF microscopy, many single molecules can be tracked over time, typically on a millisecond to minute timescale, revealing dynamic processes and structural changes.

Following this, Illumina's bridge amplification process generates clusters of identical copies from single template molecules, which are then subjected to sequencing by synthesis. This sequencing method produces a FASTQ file, a widely used format for storing sequence data alongside quality scores. In the case of Illumina sequencing, the FASTQ file also records the position of each sequencing tile and cluster positions within the tile. In the final stage, each molecule's sequence is coupled to its corresponding fluorescence by linking the sequencing position to the spatial coordinates of the single-molecule time traces, yielding a dataset containing the sequence of hundreds of thousands to millions of single-molecule fluorescence time trajectories. These data can then be analyzed using various computational and visualization approaches to elucidate sequence-dependent molecular dynamics in libraries containing up to tens of thousands of members. To facilitate adoption of MUSCLE and SPARXS,

Box1. Defining “multiplex” in single-molecule methods

The term “multiplex” in the context of single-molecule techniques can refer to distinct experimental strategies. One common usage simply denotes the parallel measurement of many individual molecules of the same type. These approaches enhance throughput and statistical resolution and are broadly applied in high-throughput single-molecule studies (122–124).

In this Review, we adopt a more specific definition: multiplex single-molecule methods that resolve and monitor multiple distinct molecular species within the same experiment. Here, multiplexing refers to molecular identity, such as differences in sequence, being discerned at the single-molecule level. These methods enable direct, comparative analyses of heterogeneous populations under identical conditions. Notably, commercial single-molecule sequencing platforms (e.g., PacBio and Oxford Nanopore) already operate in a massively parallel fashion with respect to sequence readout. However, they have generally not been used to multiplex dynamic single-molecule phenotypes in the sense defined here. Our focus is on methods that explicitly couple sequence identity to kinetic or conformational readouts across many distinct molecules in the same experiment.

standardize experimental procedures, and enhance reproducibility across laboratories, the accompanying software and test datasets have been made freely available and detailed protocol papers have been published (75, 76).

The most direct application of this approach is to investigate how DNA sequence influences conformational equilibria and dynamic behavior of structured DNA elements. For example, SPARXS has enabled the systematic analysis of the sequence dependence of the Holliday junction core across the complete sequence space (72) (Fig. 3B). Although the influence of core and flanking sequences on junction dynamics has been recognized for decades (77, 78), a quantitative high-resolution experimental study had not been made possible until recently. By applying SPARXS, the authors of this study examined all 4096 possible core sequences of the junction, generating a rich dataset that allowed them to construct a thermodynamic model capturing the underlying kinetics with high accuracy. This analysis revealed that both the core and penultimate bases contribute substantially to the dynamic behavior of the junction, a concept that had been qualitatively proposed, but not quantitatively validated, until now. Furthermore, the model enabled direct estimation of dinucleotide stacking energies, which showed strong agreement with values obtained from bulk assays. As another example involving only DNA, MUSCLE has been used to systematically characterize DNA hairpin structures using libraries of up to 4096 distinct sequences, examining their properties at several different temperatures (71).

To illustrate how this approach can be extended to protein–nucleic acid interactions, MUSCLE has also been applied to investigate the unwinding–rewinding dynamics of target DNA induced by Cas9 (71), a programmable nuclease widely used in research, biotechnology, and gene editing (79, 80) (Fig. 3C). These analyses revealed that the rate of DNA rewinding is the primary determinant of the equilibrium between unwound and rewound states. Target sequences that remained longer in the unwound conformation were more efficiently cleaved, consistent with a key role for the unwinding–rewinding equilibrium in Cas9 activity. Furthermore, by exploring a broad sequence space, several target sequences were identified that exhibited unexpected behaviors and substantially deviated from the general trend. This highlights a major advantage of comprehensive, high-throughput analyses: their independence from *a priori* assumptions, which enhances their capacity to reveal unanticipated phenomena.

SPIN-seq: Single-molecule base calling and reaction monitoring

Large-scale single-molecule profiling of DNA libraries can, however, also be achieved without cluster formation and subsequent sequencing by synthesis. Building on the ability to identify DNA sequences within short single-stranded gaps through transient binding of short fluorescent DNA oligos (44), SPIN-seq provides an additional approach for single-molecule multiplexing (73). Demonstrating the ability to work with libraries, Hazra *et al.* related the single-molecule phenotype to its corresponding DNA sequence by performing both the functional and the sequencing assays on the same solid support and instrument (Fig. 3D). The protocol starts with a DNA library featuring permutations of the sequence segment under investigation that are immobilized randomly on a glass surface. The DNA library is then interrogated for interactions with a protein of interest (or other biomolecules) using single-molecule fluorescence imaging, enabling the observation of thousands of individual molecules. These interactions also include real-time, nonequilibrium reactions driven by large multisubunit proteins. Finally, SPIN-seq is used to determine the sequence of each DNA molecule and link it to the corresponding single-molecule phenotype. The sequencing assay presently achieves high accuracy for sequences up to five nucleotides in length, thereby covering up to 1024 distinct DNA sequences. This approach enabled a detailed investigation of RNA polymerase kinetics during initial transcription (Fig. 3D), providing substantial insight into the mechanisms underlying key transitions in this complex reaction pathway.

Expanding the impact of multiplexed single-molecule studies

The ability to probe large sequence libraries with single-molecule techniques marks a major advance in molecular biology, enabling the connection of sequence identity with structural information and functional kinetics across millions of molecules in parallel. To fully realize its potential, future efforts should focus on further increasing throughput, refining approaches to study nonequilibrium enzymatic processes, and broadening applications beyond nucleic acids (Fig. 4).

Overcoming present limitations in throughput

One of the main challenges, despite highly automated implementations, remains throughput, particularly when compared with NGS platforms used in MUSCLE and SPARXS. The Illumina MiSeq, for example, can yield tens of millions of reads. However, when combined with sequential single-molecule fluorescence imaging, the yield reduces to only $\sim 10^6$ of linked molecules, owing to incomplete cluster formation, bleaching events, overlapping positions, and stringent filtering of sequencing reads and time traces. Alternatively, SPIN-seq has demonstrated high accuracy for sequences up to five nucleotides in length, thereby covering up to 1024 distinct DNA sequences. However, as sequence length increases, the number of possible sequences grows exponentially. For example, a 10-nucleotide sequence space encompasses more than 1,048,576 unique sequences. This rapid expansion makes complete sequence sampling beyond 10 nucleotides impractically time consuming, even with increased throughput.

A key strategy for increasing the efficiency of data collection is to expand the field of view (FOV) (Fig. 4A). Modern sCMOS (scientific complementary metal-oxide–semiconductor) cameras offer higher resolution than the EMCCD (electron-multiplying charge-coupled device) cameras used in early multiplexed single-molecule implementations. Using the full FOV could enhance acquisition efficiency by up to an order of magnitude. Additionally, for slower kinetic processes occurring on timescales of several seconds or more, imaging efficiency can be further increased by cycling through multiple FOVs sequentially at each time point. Acquisition efficiency can also be enhanced by methods such as DyeCycling (81) and REFRESH (82), which sustain donor fluorescence in single-molecule FRET by continuously exchanging photobleached dyes during TIRF excitation,




Fig. 3. Quantitative insights from multiplexed single-molecule platforms reveal sequence-dependent variation in kinetic and thermodynamic behavior. (A) Overview of a MUSCLE or SPARXS experiment (the example shown here is Cas9 DNA unwinding). A DNA sequence library is synthesized and immobilized on a flow cell. Single-molecule fluorescence imaging yields time traces from individual molecules. The same flow cell undergoes NGS to identify the sequence and spatial position of each DNA cluster. Fluorescence and sequencing data are aligned to extract kinetic or equilibrium parameters, such as FRET states or transitions, per sequence. [Adapted with permission from Elsevier (9); adapted with permission from (71); adapted with permission from Springer Nature (76)] (B) Sequence-dependent Holliday junction dynamics in a 4096-member SPARXS library. (i) Bases 3, 4, 7, and 8 are randomized. (ii) Heatmap of the dynamic molecule fraction across variants, with clustering seen at bases 7:8. Stars indicate fully matched junctions. (iii) Transition rates are reduced in sequences with the RYC motif. [Adapted with permission from (72)] (C) Cas9 unwinding and cleavage dynamics using MUSCLE. (i) smFRET assay with a DNA library varying four PAM-distal bases. (ii) FRET traces for a mismatched sequence (top) and equilibrium constants for all sequences (bottom), organized by identity at positions 17 to 20. (iii) Unwinding-rewinding rates plotted versus mean FRET, with model fits. (iv) FRET traces and cleavage for selected mismatched sequences show impaired cleavage correlating with FRET, linking kinetics to outcomes. [Adapted with permission from (71)] (D) SPIN-seq. (Left) Immobilized DNA libraries are imaged for biomolecular interactions (e.g., protein binding kinetics), denatured, and rehybridized. Sequence is read by hybridizing fluorescent “seals” to gapped DNA; seal binding distinguishes complementary from mismatched sequences. (Top right) Sequential interrogation of a DNA gap by four seals, with T-seal as the match. (Bottom right) smFRET monitoring of transcription initiation reveals distinct promoter DNA pathways and measures pause lifetimes across a 16-sequence library (+6/+7 degenerate bases). [Adapted from (44), CC BY 4.0; adapted from (73), CC BY 4.0]




Fig. 4. Expanding multiplexed single-molecule analysis across molecular scales and applications. (A) Increasing data collection efficiency. Throughput can be maximized by (i) using full CMOS camera fields of view or (ii) cycling through multiple FOVs sequentially for slower kinetic processes. (iii) DyeCycling (81) sustains donor fluorescence in single-molecule FRET by continuously exchanging photobleached dyes during TIRF excitation, enabling longer observation windows for kinetic analysis and increasing the statistical power of equilibrium binding measurements from fewer molecules. (B) Peptide and protein libraries. DNA-barcoded peptide and protein libraries allow single-molecule kinetic screening of binding, folding, or enzymatic activity. Techniques such as ribosome and cDNA display link behavior to sequence, enabling functional profiling across millions of variants. Applications span signaling, immunity, and protein engineering. (C) Aptamer characterization. High-throughput screening of nucleic acid aptamers at the single-molecule level can uncover folding pathways, binding kinetics, and structural dynamics tied to sequence. This enables identification of aptamers with high specificity, adaptability, and functional stability for therapeutic, diagnostic, and analytical uses. (D) Sequence multiplexing in other single-molecule methods, such as magnetic tweezers. Incorporating sequence-encoded tethers enables parallel force spectroscopy on multiple DNA or RNA constructs, facilitating comparison of mechanical responses or enzymatic processing. These platforms can also include colocalization-based fluorescence readouts for monitoring protein dynamics or substrate conformational changes alongside mechanical data. (E) Expansion to cell libraries. Linking single-molecule readouts to cell-based assays involves using genetically defined cell libraries combined with phenotypic sorting (e.g., growth, stress). Cells can be analyzed for molecular structure, localization, and dynamics using FRET, MINFLUX (128), expansion microscopy (129), and smFISH (130,131), connecting genotype, phenotype, and mechanism in a unified experimental pipeline. Visual examples are stylized illustrations inspired by reported single-molecule observations in cells.

thereby enabling longer observation windows for kinetic analysis. Advances in hardware and software automation, alongside optimized imaging protocols to maximize the fraction of molecules that can be successfully linked, will be critical in improving the accessibility, efficiency, and throughput of multiplexed single-molecule studies.

It is important to note, however, that the strength of multiplexed single-molecule approaches lies not in replacing high-throughput sequencing but in complementing it by providing kinetic and mechanistic insights that bulk sequencing cannot capture. As such, single-molecule sequence profiling may be best suited for targeted studies, focusing on biologically relevant sequences rather than random libraries. A genome-referenced approach, as demonstrated in loop-seq (2) provides a promising strategy for prioritizing relevant sequences. By anchoring analysis to a well-defined reference framework, such as the yeast genome or specific regulatory elements, kinetic parameters from tens of thousands of sequences involved in key processes such as transcription, translation, and chromatin dynamics can be efficiently extracted.

When it is impractical to exhaustively sample the vast sequence space, machine learning (ML) offers a powerful tool for extracting meaningful insights from incomplete datasets. ML can identify patterns, classify molecular behaviors, and infer missing kinetic parameters from sparsely sampled data. Nonetheless, broad sequence scans remain essential, as they often uncover rare behaviors that ML models trained on limited data may miss. Recent work shows that large language models trained on biological sequences and kinetic measurements can capture rich sequence-function relationships (83), suggesting that the high-throughput datasets produced by MUSCLE, SPARXS, and SPIN-seq could serve as particularly powerful training material for such models. Conversely, single-molecule sequence profiling can serve as an experimental platform to validate ML-predicted interaction kinetics and binding specificities. The iterative feedback between data-driven prediction, continuous experimental testing, and refined modeling will ultimately enhance the interpretability and utility of these technologies.

Advancing methodology: Nonequilibrium studies

Many fundamental biological processes, such as transcription, translation, and ligand binding, occur under nonequilibrium conditions and involve transient molecular states that are difficult to capture using ensemble methods. A key challenge is the study of nucleoside triphosphate (NTP)-hydrolyzing enzymes, such as helicases and polymerases, which drive irreversible modifications of biological macromolecules. In single-molecule experiments, these nonequilibrium processes are typically initiated by perfusing the flow cell with enzymes and/or NTP cofactors. Only one FOV can be imaged because, after data acquisition, the reaction has already occurred across the entire surface of the flow cell. Consequently, only a small fraction of the substrates can be observed. To overcome this issue, photo-uncaging strategies such as LAGOON (84) provide spatiotemporal control over enzymatic reactions, enabling precise synchronization between reaction initiation and imaging. Integrating LAGOON or other light-based reaction control strategies

(85–87) will therefore enable multiplexed single-molecule observations of complex nonequilibrium dynamics.

Broadening applications

Although multiplexed single-molecule studies have primarily been applied to DNA libraries and can be readily adapted to RNA, its methodology can also be extended to protein libraries, thereby broadening its impact in molecular biology (Fig. 4B). A key application is the study of protein-protein and peptide-protein interactions, which play a crucial role in critical processes such as signal transduction and immune recognition. By incorporating cDNA display or ribosome display for peptide and protein libraries (88, 89), MUSCLE, SPARXS, or SPIN-seq could enable high-throughput kinetic analyses of millions of individual peptides and proteins in parallel. Recent advances have paved the way for such approaches, including the profiling of DNA-barcoded protein interactions at the single-molecule level (90), real-time kinetic screening of thousands of protein interactions using multiplexed biosensors (91), and the integration of ribosome display with deep sequencing for rapid antibody discovery (92).

Multiplexed single-molecule profiling also holds promise for investigating protein-ligand interactions, which play central roles in receptor signaling and drug binding (Fig. 4C). A notable example is the use of nucleic acid aptamers—short, structured DNA or RNA molecules selected for high-affinity binding to specific protein targets—which can be integrated into libraries for parallel interaction profiling. Traditional high-throughput screening methods rely primarily on affinity-based measurements, which fail to capture critical kinetic parameters, such as association and dissociation rates, which are key determinants of drug efficacy and specificity. Multiplexed single-molecule studies offer an alternative, enabling real-time kinetic screening of ligand-target interactions across large target and chemical libraries. For example, although DNA-encoded chemical libraries (93–95) contain millions to billions of small molecules, their integration with single-molecule kinetic measurements remains an unexplored frontier. Bridging this gap could allow multiplexed single-molecule approaches to identify compounds with optimal kinetic profiles for therapeutic development.

Although the MUSCLE, SPARXS, and SPIN-seq approaches thus far have been used primarily for single-molecule FRET (smFRET) experiments, they are also compatible with other single-molecule fluorescence techniques, including colocalization analysis and protein-induced fluorescence enhancement (96). Furthermore, single-molecule localization microscopy methods can be integrated, such as PALM, STORM, and DNA-PAINT (97–99), enabling the exploration of how sequence variations influence nano- and microscale structural organization. Many biological processes involve structural transitions that span hundreds of nanometers to micrometers. Investigating such large-scale dynamics often requires techniques capable of probing molecular mechanics over extended distances while maintaining single-molecule resolution. Integrating force-spectroscopy methods, particularly magnetic tweezers (100–103), into the MUSCLE, SPARXS, and SPIN-seq approaches would enable high-throughput studies of these long-range biomolecular interactions (Fig. 4D). Additionally, large rigid DNA scaffolds constructed using DNA origami (104–107) can be readily combined with oligonucleotide libraries and integrated, enabling the study of large-scale dynamics beyond the spatial range accessible by smFRET. Building on the origami-rotor-based imaging and tracking (ORBIT) approach (108), DNA origami can also be used within multiplexed single-molecule studies to access a qualitatively different mode of motion—DNA rotation—that is not readily detectable by smFRET. However, overly large structures may interfere with cluster or gap formation, and experimental validation of these approaches may reveal additional practical limitations.

Although the MUSCLE, SPARXS, and SPIN-seq approaches enable the direct linkage of complex molecular dynamics to sequence identity,

they are inherently *in vitro* based and may not fully capture the dynamic behaviors that occur within native cellular environments. One strategy to bridge this gap involves the use of cellular extracts applied to immobilized nucleic acid libraries, thereby recreating more physiologically relevant conditions. Additionally, the emergence of *in situ* genotyping and sequencing techniques, such as rolling circle amplification, barcoding strategies, and related methods (109–112), has opened new avenues for coupling genotypic information with multiplexed phenotypic readouts. Proof-of-concept studies for optical pooled screening in both bacterial and mammalian systems have already demonstrated the feasibility of combining live-cell phenotyping with *in situ* genotyping or barcoded readouts (113–116).

Naturally, the *in vivo* environment presents major experimental challenges, including limited spatiotemporal resolution, lower signal-to-noise ratios, and constraints on the labeling and detection of molecular species (117–121). Substantial technological advances will be required to bring *in vivo* single-molecule approaches closer to their *in vitro* counterparts in terms of data quality. However, such developments may one day provide similarly granular mechanistic insights within native cellular contexts (Fig. 4E).

A powerful reunion of sequencing and single-molecule biophysics

Taken together, the developments discussed here underscore the ongoing convergence of two foundational technologies in modern biology: sequencing and single-molecule biophysics. By fusing the throughput and scalability of advanced sequencing technologies with the mechanistic and dynamic insight of single-molecule imaging, we are now poised to interrogate complex biological phenomena with previously unattainable depth and precision. Whether on a flow cell or within a living cell, the ability to link molecular identity with dynamic behavior, across millions of distinct molecules, ushers in a new era of discovery. This represents the latest chapter in a recurring narrative, where sequencing and single-molecule biophysics repeatedly intersect to advance our understanding of life's molecular underpinnings.

REFERENCES AND NOTES

1. E. Oberbeckmann *et al.*, *Nat. Commun.* **12**, 3231 (2021).
2. A. Basu *et al.*, *Nature* **589**, 462–467 (2021).
3. J. M. Tome *et al.*, *Nat. Methods* **11**, 683–688 (2014).
4. R. Nutiu *et al.*, *Nat. Biotechnol.* **29**, 659–664 (2011).
5. C. Jung *et al.*, *Cell* **170**, 35–47.e13 (2017).
6. R. E. Kleiner, C. E. Dumelin, G. C. Tiu, K. Sakurai, D. R. Liu, *J. Am. Chem. Soc.* **132**, 11779–11791 (2010).
7. U. T. Nguyen *et al.*, *Nat. Methods* **11**, 834–840 (2014).
8. G. P. Dann *et al.*, *Nature* **548**, 607–611 (2017).
9. I. Severins, C. Joo, J. van Noort, *Mol. Cell* **82**, 1788–1805 (2022).
10. J. B. Kinney, D. M. McCandlish, *Annu. Rev. Genomics Hum. Genet.* **20**, 99–127 (2019).
11. E. Marklund, Y. Ke, W. J. Greenleaf, *Nat. Rev. Genet.* **24**, 401–414 (2023).
12. C. J. Layton, P. L. McMahon, W. J. Greenleaf, *Mol. Cell* **73**, 1075–1082.e4 (2019).
13. E. A. Boyle *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **114**, 5461–5466 (2017).
14. N. Svensen, O. B. Peersen, S. R. Jaffrey, *Chembiochem* **17**, 1628–1635 (2016).
15. L. C. Szymczak, H.-Y. Kuo, M. Mrksich, *Anal. Chem.* **90**, 266–282 (2018).
16. D. Wu *et al.*, *Anal. Chem.* **95**, 2645–2652 (2023).
17. Z. Li *et al.*, *Sci. Adv.* **6**, eabb3350 (2020).
18. N. Mamet *et al.*, *bioRxiv* 630830 [Preprint] (2019); <https://doi.org/10.1101/630830>.
19. J. D. Buenrostro *et al.*, *Nat. Biotechnol.* **32**, 562–568 (2014).
20. A. Melnikov *et al.*, *Nat. Biotechnol.* **30**, 271–277 (2012).
21. R. Tewhey *et al.*, *Cell* **165**, 1519–1529 (2016).
22. D. M. Fowler *et al.*, *Nat. Methods* **7**, 741–746 (2010).
23. C. L. Araya, D. M. Fowler, *Trends Biotechnol.* **29**, 435–442 (2011).
24. A. Jolma *et al.*, *Cell* **152**, 327–339 (2013).
25. S. Mohapatra, C.-T. Lin, X. A. Feng, A. Basu, T. Ha, *Chem. Rev.* **120**, 36–78 (2020).
26. L. Bacic, A. Sabantsev, S. Deindl, *Curr. Opin. Struct. Biol.* **65**, 61–68 (2020).
27. F. R. Hill, E. Monachino, A. M. van Oijen, *Biochem. Soc. Trans.* **45**, 759–769 (2017).
28. J. Choi, R. Grosely, E. V. Puglisi, J. D. Puglisi, *Curr. Opin. Struct. Biol.* **58**, 233–240 (2019).
29. M. Orrit, T. Ha, V. Sandoghdar, *Chem. Soc. Rev.* **43**, 973–976 (2014).
30. F. Kulzer, M. Orrit, *Annu. Rev. Phys. Chem.* **55**, 585–611 (2004).

31. M. Dangkulwanich, T. Ishibashi, L. Bintu, C. Bustamante, *Chem. Rev.* **114**, 3203–3223 (2014).
32. J. Zhou, V. Schweikhard, S. M. Block, *Biochim. Biophys. Acta* **1829**, 29–38 (2013).
33. J. Michaelis, B. Treutlein, *Chem. Rev.* **113**, 8377–8399 (2013).
34. L. Bai, T. J. Santangelo, M. D. Wang, *Annu. Rev. Biophys. Biomol. Struct.* **35**, 343–360 (2006).
35. M. Stracy, A. N. Kapanidis, *Methods* **120**, 103–114 (2017).
36. E. Lerner *et al.*, *Science* **359**, eaan1133 (2018).
37. D. Dulin, B. A. Berghuis, M. Depken, N. H. Dekker, *Curr. Opin. Struct. Biol.* **34**, 116–122 (2015).
38. M. F. Juetten *et al.*, *Curr. Opin. Chem. Biol.* **20**, 103–111 (2014).
39. J. H. Felce, S. J. Davis, D. Klenerman, *Trends Pharmacol. Sci.* **39**, 96–108 (2018).
40. J. Camunas-Soler, M. Ribezzi-Crivellari, F. Ritort, *Annu. Rev. Biophys.* **45**, 65–84 (2016).
41. S. Park, T. Ha, G. D. Bowman, *Nucleic Acids Res.* **51**, 10326–10343 (2023).
42. A. A. Hoskins, J. Gelles, M. J. Moore, *Curr. Opin. Chem. Biol.* **15**, 864–870 (2011).
43. S. Deindl, X. Zhuang, *Methods Enzymol.* **513**, 59–86 (2012).
44. R. Andrews *et al.*, bioRxiv 2022.02.27482175 [Preprint] (2022); <https://doi.org/10.1101/2022.02.27482175>.
45. K. Makasheva *et al.*, *J. Am. Chem. Soc.* **143**, 16313–16319 (2021).
46. J. Chu *et al.*, *Nat. Nanotechnol.* **19**, 1150–1157 (2024).
47. E. Shema *et al.*, *Science* **352**, 717–721 (2016).
48. V. Fedyuk *et al.*, *Nat. Biotechnol.* **41**, 212–221 (2023).
49. L. M. Davis *et al.*, *Genet. Anal. Tech. Appl.* **8**, 1–7 (1991).
50. I. Braslavsky, B. Hebert, E. Kartalov, S. R. Quake, *Proc. Natl. Acad. Sci. U.S.A.* **100**, 3960–3964 (2003).
51. M. J. Levene *et al.*, *Science* **299**, 682–686 (2003).
52. W. J. Greenleaf, S. M. Block, *Science* **313**, 801 (2006).
53. T. D. Harris *et al.*, *Science* **320**, 106–109 (2008).
54. J. Eid *et al.*, *Science* **323**, 133–138 (2009).
55. J. Clarke *et al.*, *Nat. Nanotechnol.* **4**, 265–270 (2009).
56. T. Ha *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **93**, 6264–6268 (1996).
57. X. Zhuang *et al.*, *Science* **288**, 2048–2051 (2000).
58. J. Liphardt, B. Onoa, S. B. Smith, I. Tinoco Jr., C. Bustamante, *Science* **292**, 733–737 (2001).
59. G. Bonnet, O. Krichavsky, A. Libchaber, *Proc. Natl. Acad. Sci. U.S.A.* **95**, 8602–8606 (1998).
60. M. D. Wang *et al.*, *Science* **282**, 902–907 (1998).
61. K. C. Neuman, E. A. Abbondanzieri, R. Landick, J. Gelles, S. M. Block, *Cell* **115**, 437–447 (2003).
62. M. H. Larson *et al.*, *Science* **344**, 1042–1047 (2014).
63. I. O. Vvedenskaya *et al.*, *Science* **344**, 1285–1289 (2014).
64. I. C. Nova *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **121**, e2321017121 (2024).
65. J. M. Craig *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **114**, 11932–11937 (2017).
66. J. M. Craig *et al.*, *Nucleic Acids Res.* **50**, 10601–10613 (2022).
67. A. H. Laszlo *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2202489119 (2022).
68. M. MacKenzie, C. Argyropoulos, *Micromachines* **14**, 459 (2023).
69. M. Palla *et al.*, *ACS Nano* **15**, 489–502 (2021).
70. S. Uemura *et al.*, *Nature* **464**, 1012–1017 (2010).
71. J. Aguirre Rivera *et al.*, *Science* **385**, 892–898 (2024).
72. I. Severins *et al.*, *Science* **385**, 898–904 (2024).
73. J. P. Hazra *et al.*, bioRxiv 2025.11.27690701 [Preprint] (2025); <https://doi.org/10.1101/2025.11.27690701>.
74. D. R. Bentley *et al.*, *Nature* **456**, 53–59 (2008).
75. C. Bastiaanssen, I. Severins, J. van Noort, C. Joo, *Nat. Protoc.* 10.1038/s41596-025-01196-y (2025).
76. M. Panfilov *et al.*, *Nat. Protoc.* 10.1038/s41596-025-01198-w (2025).
77. S. M. Miick, R. S. Fee, D. P. Millar, W. J. Chazin, *Proc. Natl. Acad. Sci. U.S.A.* **94**, 9080–9084 (1997).
78. D. M. Lilley, R. M. Clegg, *Q. Rev. Biophys.* **26**, 131–175 (1993).
79. M. Jinek *et al.*, *Science* **337**, 816–821 (2012).
80. A. V. Anzalone, L. W. Koblan, D. R. Liu, *Nat. Biotechnol.* **38**, 824–844 (2020).
81. B. Vermeer, S. Schmid, *Nano Res.* **15**, 9818–9830 (2022).
82. M. Kümmerlin, A. Mazumder, A. N. Kapanidis, *Chemphyschem* **24**, e202300175 (2023).
83. G. Wei, X. Ran, R. Ai-Absi, Z. Yang, *Protein Sci.* **34**, e70251 (2025).
84. A. Sabantsev *et al.*, *Nat. Chem. Biol.* **18**, 1144–1151 (2022).
85. A. Korman *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **117**, 12080–12086 (2020).
86. C. Brieke, F. Rohrbach, A. Gottschalk, G. Mayer, A. Heckel, *Angew. Chem. Int. Ed.* **51**, 8446–8476 (2012).
87. Y. Liu *et al.*, *Science* **368**, 1265–1269 (2020).
88. J. Yamaguchi *et al.*, *Nucleic Acids Res.* **37**, e108 (2009).
89. J. Hanes, A. Plückthun, *Proc. Natl. Acad. Sci. U.S.A.* **94**, 4937–4942 (1997).
90. L. Gu *et al.*, *Nature* **515**, 554–557 (2014).
91. C. V. Agu *et al.*, *Commun. Biol.* **8**, 468 (2025).
92. B. T. Porebski *et al.*, *Nat. Biomed. Eng.* **8**, 214–232 (2024).
93. M. C. Needels *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **90**, 10700–10704 (1993).
94. S. Melkko, J. Scheuermann, C. E. Dumelin, D. Neri, *Nat. Biotechnol.* **22**, 568–574 (2004).
95. S. Brenner, R. A. Lerner, *Proc. Natl. Acad. Sci. U.S.A.* **89**, 5381–5383 (1992).
96. H. Hwang, H. Kim, S. Myong, *Proc. Natl. Acad. Sci. U.S.A.* **108**, 7414–7418 (2011).
97. E. Betzig *et al.*, *Science* **313**, 1642–1645 (2006).
98. M. J. Rust, M. Bates, X. Zhuang, *Nat. Methods* **3**, 793–796 (2006).
99. R. Jungmann *et al.*, *Nat. Methods* **11**, 313–318 (2014).
100. S. B. Smith, L. Finzi, C. Bustamante, *Science* **258**, 1122–1126 (1992).
101. T. R. Strick, J. F. Allemand, D. Bensimon, A. Bensimon, V. Croquette, *Science* **271**, 1835–1837 (1996).
102. J. P. Cossens, D. Dulin, N. H. Dekker, *Rev. Sci. Instrum.* **85**, 103712 (2014).
103. A. Huhle *et al.*, *Nat. Commun.* **6**, 5885 (2015).
104. P. W. K. Rothmund, *Nature* **440**, 297–302 (2006).
105. S. M. Douglas *et al.*, *Nature* **459**, 414–418 (2009).
106. D. Han *et al.*, *Science* **332**, 342–346 (2011).
107. N. C. Seeman, *J. Theor. Biol.* **99**, 237–247 (1982).
108. P. Kosuri, B. D. Altheimer, M. Dai, P. Yin, X. Zhuang, *Nature* **572**, 136–140 (2019).
109. C. Larsson, I. Grundberg, O. Söderberg, M. Nilsson, *Nat. Methods* **7**, 395–397 (2010).
110. R. Ke *et al.*, *Nat. Methods* **10**, 857–860 (2013).
111. J. H. Lee *et al.*, *Science* **343**, 1360–1363 (2014).
112. C. Xia, J. Fan, G. Emanuel, J. Hao, X. Zhuang, *Proc. Natl. Acad. Sci. U.S.A.* **116**, 19490–19499 (2019).
113. M. J. Lawson *et al.*, *Mol. Syst. Biol.* **13**, 947 (2017).
114. G. Emanuel, J. R. Moffitt, X. Zhuang, *Nat. Methods* **14**, 1159–1162 (2017).
115. A. Farrar *et al.*, *Commun. Biol.* **8**, 319 (2025).
116. D. Feldman *et al.*, *Cell* **179**, 787–799.e17 (2019).
117. A. Gahlmann, W. E. Moerner, *Nat. Rev. Microbiol.* **12**, 9–22 (2014).
118. J. Elf, I. Barkefors, *Annu. Rev. Biochem.* **88**, 635–659 (2019).
119. M. Lelek *et al.*, *Nat. Rev. Methods Primers* **1**, 39 (2021).
120. N. K. Podh *et al.*, *J. Mol. Biol.* **433**, 167250 (2021).
121. T. Ha, C. Kaiser, S. Myong, B. Wu, J. Xiao, *Mol. Cell* **82**, 304–314 (2022).
122. N. Ribbeck, O. A. Saleh, *Rev. Sci. Instrum.* **79**, 094301 (2008).
123. D. Yang, A. Ward, K. Halvorsen, W. P. Wong, *Nat. Commun.* **7**, 11026 (2016).
124. R. Agarwal, K. E. Duderstadt, *Nat. Commun.* **11**, 4714 (2020).
125. D. Huynh *et al.*, *Nat. Commun.* **16**, 1259 (2025).
126. H. Shi *et al.*, *Mol. Cell* **85**, 1730–1742.e9 (2025).
127. E. A. Abbondanzieri, W. J. Greenleaf, J. W. Shaevitz, R. Landick, S. M. Block, *Nature* **438**, 460–465 (2005).
128. F. Balzarotti *et al.*, *Science* **355**, 606–612 (2017).
129. F. Chen, P. W. Tillberg, E. S. Boyden, *Science* **347**, 543–548 (2015).
130. A. M. Femino, F. S. Fay, K. Fogarty, R. H. Singer, *Science* **280**, 585–590 (1998).
131. A. Raj, P. van den Bogaard, S. A. Rifkin, A. van Oudenaarden, S. Tyagi, *Nat. Methods* **5**, 877–879 (2008).

ACKNOWLEDGMENTS

We thank M. Panfilov and A. Sabantsev for critical reading of the manuscript. **Funding:** This work was funded by European Research Council (ERC) Advanced Grant ERC-ADG-101092623 (S.D.); Knut and Alice Wallenberg Foundation grants KAW/WAF 2019.0306 and KAW 2024.0012 (S.D.); Cancerfonden grant 25 4453 Pj (S.D.); Swedish Research Council project grant VR 03255 (S.D.); ERC Consolidator Grant 819299 (C.J.); Bio&Medical Technology Development Program of the National Research Foundation RS-2025-02217909 (C.J.) and the Basic Research Laboratory Program NRF-2023R1A2C2004745 (C.J.) by the Ministry of Science and ICT, Korea; Wellcome Trust grants 110164/Z/15/Z and 226662/Z/22/Z (A.N.K.); and Leverhulme Trust award grant RPG-2024-037 (A.N.K., J.P.H.). **Competing interests:** This work discusses single-molecule fluorescence measurements, which can be performed using commercial microscopes from Oxford Nanosystems, a company in which A.N.K. is a co-founder and shareholder. J.P.H. and A.N.K. have financial interests in patent applications related to the SPIN-seq methodology discussed in this manuscript. **License information:** Copyright © 2026 the authors, some rights reserved; exclusive license American Association for the Advancement of Science. No claim to original US government works. <https://www.science.org/about/science-licenses-journal-article-reuse>

Submitted 4 June 2025; accepted 9 December 2025

10.1126/science.adv4503



From sequence to function: Bridging single-molecule kinetics and molecular diversity

A. N. Kapanidis, L. Muras, K. Sreenivasa, J. P. Hazra, J. van Noort, C. Joo, and S. Deindl

Science **391** (6784), . DOI: 10.1126/science.adv4503

Editor's summary

How DNA and RNA molecules fold, move, and interact is controlled by their sequence in complex ways that remain obscure. Existing methods can either determine sequences or measure single-molecule dynamics, but they typically do not connect both types of information for large numbers of distinct sequences at once. Kapanidis *et al.* reviewed new approaches that track the real-time dynamics of hundreds of thousands of individual molecules and then read out the exact sequence of each one. By linking sequence to kinetic “behavior profiles,” these methods open new routes to understanding mutations, drug action, and molecular mechanisms. —Di Jiang

View the article online

<https://www.science.org/doi/10.1126/science.adv4503>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)

Science (ISSN 1095-9203) is published by the American Association for the Advancement of Science. 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2026 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works