

## Surface melt on Antarctic ice shelves driven by wind-albedo interactions

Lhermitte, Stef; Lenaerts, Jan T M

**Publication date**

2017

**Document Version**

Final published version

**Published in**

Geophysical Research Abstracts (online)

**Citation (APA)**

Lhermitte, S., & Lenaerts, J. T. M. (2017). Surface melt on Antarctic ice shelves driven by wind-albedo interactions. *Geophysical Research Abstracts (online)*, 19, Article EGU2017-13779.

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.



## **Surface melt on Antarctic ice shelves driven by wind-albedo interactions**

Stef Lhermitte (1) and Jan Lenaerts (2)

(1) TUDelft, Department of Geoscience & Remote Sensing, Netherlands (s.lhermitte@tudelft.nl), (2) Utrecht University, Institute for Marine and Atmospheric Research Utrecht, The Netherlands.

Surface melt and subsequent firn air depletion is considered an important precursor for disintegration of Antarctic ice shelves, causing grounded glaciers to accelerate and sea level to rise. Recent studies have highlighted the impact of surface winds on Antarctic ice shelf melt, both on the Antarctic Peninsula and in East Antarctica. In the Antarctic Peninsula, foehn winds enhance melting near the grounding line, which in the recent past has led to the disintegration of the most northerly ice shelves. On the East Antarctic ice shelves, on the other hand, meltwater-induced firn air depletion is found in the grounding zone as result of persistent katabatic winds, regionally warming the atmosphere and inducing a melt-albedo feedback.

Here, we use a combination multi-source satellite imagery, snow modelling, climate model output and in-situ observations to highlight the importance of this wind-induced melt and to show its widespread occurrence across Antarctica. The satellite imagery gives insight in the meltwater drainage systems, showing spatio-temporal changes in both supraglacial and englacial water throughout the melt season and during the subsequent winter.

Although the wind-induced melt is a regional phenomenon with strong inter-annual variability, it is strongly correlated to larger scale climate parameters, such as summer surface temperature. Based on these correlations and snow model output driven by future climate scenarios, we can constrain the future changes to this local melt near the grounding line.