

## The Markerwadden

### the influence of the sand fraction on the ripening behaviour of Markermeer sediment

Barciela Rial, Maria; Winterwerp, Han; van Paassen, Leon; Griffioen, Jasper; van Kessel, Thijs

**Publication date**

2016

**Document Version**

Final published version

**Citation (APA)**

Barciela Rial, M., Winterwerp, H., van Paassen, L., Griffioen, J., & van Kessel, T. (2016). *The Markerwadden: the influence of the sand fraction on the ripening behaviour of Markermeer sediment*. 10-10. Abstract from NCK-days 2016, Ouddorp, Netherlands.

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.

# **THE MARKERWADDEN: THE INFLUENCE OF THE SAND FRACTION ON THE RIPENING BEHAVIOUR OF MARKERMEER SEDIMENT**

Maria Barciela Rial<sup>1\*</sup>, Johan C. Winterwerp<sup>1,2</sup>, Leon A. van Paassen<sup>1</sup>, Jasper Griffioen<sup>3,4</sup> and Thijs van Kessel<sup>2</sup>

<sup>1</sup> Delft University of Technology, Faculty of Civil Engineering and Geosciences -\*m.barcielarial@tudelft.nl

<sup>2</sup> Deltares PO Box 177, 2600 MH Delft, The Netherlands

<sup>3</sup> Utrecht University, Department of Innovation, Environmental and Energy Sciences, Faculty of Geosciences.

<sup>4</sup> TNO P.O. Box 80015, 3508 TA Utrecht, The Netherlands

## **Description of research**

The large and shallow Lake *Markermeer*, The Netherlands suffers from ecological problems. Its ecosystem has been affected by high turbidity levels, caused by the resuspension of the oxic and highly bioturbated uppermost layer of the bed. To improve its water quality, a broad and multidisciplinary consortium has been working together since 2013 on the *MarkerWadden*. This Building with Nature (BwN) project aims to improve the ecosystem in the lake by creating a wetland with sediments partly originating from the mud material of the bed of the lake itself. Part of the sediments which is placed above the water level will be exposed to evaporative conditions. Due to evaporation these sediments will consolidate. The deformation behaviour of ripening sediments is significantly affected by its composition. The sediment composition varies enormously in the *Markermeer*, in particular the sand content. In this research the behaviour of sediment while drying was carried out.

Figure 1 shows a fragment of the negative excess pore pressure development in time at the bottom probe of a Hyprop for a natural *Markemeer* Holocene clay sample (containing 10% of sand) and a sieved sample (without sand) during a desiccation test in a climate room with a constant temperature of 24°C.




Figure 1. Negative excess pore pressure at the bottom probe corrected for the initial hydrostatic head and the hydrostatic head loss in time as a result of evaporation and shrinkage

From the water loss and measured pore pressures the water retention curves are determined (Figure 2a). The curves have a different starting point and clearly do not align at higher suction values due to a difference in the initial liquidity and a different composition. Figure 2b shows that the undrained shear strength as a function of liquidity index of the remoulded natural sample clearly fits the empirical correlations available in literature.




Figure 2. a) Water retention curve of a sieved and a natural sample of Markermeer Holocene clay. b) Correlation between the liquidity index and the undrained remoulded shear strength for a natural sample