

## Assessing the climate impact of formation flights

Frömming, C.; Grewe, V.; Brinkop, S.; Haslerud, Amund S.; Rosanka, S.; Matthes, Sigrun ; van Manen, J.

**Publication date**  
2020

**Document Version**  
Final published version

**Citation (APA)**  
Frömming, C., Grewe, V., Brinkop, S., Haslerud, A. S., Rosanka, S., Matthes, S., & van Manen, J. (2020). *Assessing the climate impact of formation flights*. 179-179. Abstract from 3rd ECATS conference.

**Important note**  
To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**  
Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**  
Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.

## WEATHER AND LOCATION DEPENDENCY OF AVIATION CLIMATE EFFECTS: 4-D-CLIMATE-CHANGE-FUNCTIONS

C. Frömming<sup>1</sup>, V. Grewe<sup>1,2</sup>, S. Brinkop<sup>1</sup>, A. S. Haslerud<sup>3</sup>, S. Rosanka<sup>1,2,4</sup>, J. van Manen<sup>2</sup> & S. Matthes<sup>1</sup>

<sup>1</sup> *Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany*

<sup>2</sup> *Delft University of Technology, Aerospace Engineering, The Netherlands*

<sup>3</sup> *Center for International Climate and Environmental Research, Oslo, Norway*

<sup>4</sup> *Forschungszentrum Jülich GmbH, Institute of Energy and Climate Research, IEK-8: Troposphere, Jülich, Germany*

**Abstract.** Emissions of aviation include CO<sub>2</sub>, H<sub>2</sub>O, NO<sub>x</sub> and particles. While CO<sub>2</sub> has a long atmospheric residence time and is uniformly distributed in the atmosphere, non-CO<sub>2</sub> gases, particles and their products have short atmospheric residence times and are heterogeneously distributed. Their climate effects depend on chemical and meteorological background conditions during emission, which vary with geographic location, altitude, time, local insolation, actual weather, etc. This spatial and temporal variability can be utilized for aviation climate impact mitigation by identifying aircraft trajectories which avoid climate-sensitive regions. To determine the climate change contribution of individual emissions as function of 3-dimensional position, time and weather situation, contributions of local emissions to changes in O<sub>3</sub>, CH<sub>4</sub>, H<sub>2</sub>O and contrail-cirrus were computed by means of the ECHAM5/MESSy Atmospheric Chemistry model and four-dimensional climate change functions (CCFs) were derived thereof. Typical weather situations in the North Atlantic region were considered for winter and summer. For all non-CO<sub>2</sub> species included in the study, we found distinct weather related differences with respect to their climate impact. Depending on the species, we found enhanced significance of the position of emission release in relation to high pressure systems, in relation to the jet stream, in relation to polar night and in relation to the tropopause altitude. The dominating parameters were found to be contrail-cirrus and total NO<sub>x</sub>. The results of this study represent a comprehensive basis for weather dependent flight trajectory optimization studies. Furthermore it constitutes the groundwork for the development of more generally applicable algorithmic CCFs.

**Keywords:** non-CO<sub>2</sub> emissions, weather dependency, climate optimal trajectories