

Evolution of Gaussian Concentration Bounds under Diffusions

Chazottes, J. R.; Collet, P.; Redig, F.

Publication date

2021

Document Version

Final published version

Published in

Markov Processes and Related Fields

Citation (APA)

Chazottes, J. R., Collet, P., & Redig, F. (2021). Evolution of Gaussian Concentration Bounds under Diffusions. *Markov Processes and Related Fields*, 27(5), 707-754.

Important note

To cite this publication, please use the final published version (if applicable). Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights. We will remove access to the work immediately and investigate your claim.



HAL
open science

Evolution of Gaussian concentration bounds under diffusions

J.-R Chazottes, P. Collet, F Redig

► **To cite this version:**

J.-R Chazottes, P. Collet, F Redig. Evolution of Gaussian concentration bounds under diffusions. Markov Processes And Related Fields, 2021, 27 (5). hal-02167399v2

HAL Id: hal-02167399

<https://hal.science/hal-02167399v2>

Submitted on 15 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evolution of Gaussian Concentration

J.-R. Chazottes ^{*1}, P. Collet ^{†1}, and F. Redig ^{‡2}

¹Centre de Physique Théorique, CNRS, Institut Polytechnique de Paris, F-91128 Palaiseau Cedex, France

²Delft Institute of Applied Mathematics, Delft University of Technology, van Mourik Broekmanweg 6, 2628 XE, Delft, The Netherlands

July 15, 2020

Abstract

We study the behavior of the Gaussian concentration bound (GCB) under stochastic time evolution. More precisely, in the context of Markovian diffusion processes on \mathbb{R}^d we prove in various settings that if we start the process from an initial probability measure satisfying GCB, then at later times GCB holds, and estimates for the constant are provided. Under additional conditions, we show that GCB holds for the unique invariant measure. This gives a semigroup interpolation method to prove Gaussian concentration for measures which are not available in explicit form. We also consider diffusions “coming down from infinity” for which we show that, from any starting measure, at positive times, GCB holds. Finally we consider non-Markovian diffusion processes with drift of Ornstein-Uhlenbeck type, and general bounded predictable variance.

Key-words: Markov diffusions, Ornstein-Uhlenbeck process, nonlinear semigroup, coupling, Bakry-Emery criterion, non-reversible diffusions, diffusions coming down from infinity, Ginzburg-Landau diffusions, non-Markovian diffusions, Lorenz attractor with noise, Burkholder inequality .

*Email: chazottes@cpht.polytechnique.fr

†Email: collet@cpht.polytechnique.fr

‡Email: f.h.j.redig@tudelft.nl

Contents

1	Introduction	2
2	Setting and basic questions	4
2.1	Gaussian concentration bounds	4
2.2	Time evolved Gaussian concentration bound	5
2.3	Illustrative example: an Ornstein-Uhlenbeck process	6
3	Nonlinear semigroup approach	7
3.1	The nonlinear semigroup	7
3.2	Some preparatory computations	8
3.3	Abstract gradient bound approach	10
4	Coupling approach	12
4.1	Coupling and the nonlinear semigroup	12
4.2	Examples	17
5	Distance Gaussian moment approach	19
5.1	A general equivalence	19
5.2	Example 1: Diffusions coming down from infinity	21
5.3	Example 2: Markovian diffusion processes with space-time dependent drift and covariance	24
6	Non Markovian diffusions: Martingale moment approach	27
A	Proof of Theorem 5.1	30
B	An approximation lemma	32

1 Introduction

Concentration inequalities are a well studied subject in probability and statistics and are very useful in the study of fluctuations of possibly complicated and indirectly defined functions of random variables, such as the Kantorovich distance between the empirical distribution and the true distribution, and various properties of random graphs. See for instance [3, 12] and references therein. Initially mostly studied in the context of independent random variables, many efforts have been done to extend concentration inequalities to the context of dependent random variables, and more generally dependent

random fields. For instance, in the context of models of statistical mechanics, where the dependence is naturally encoded in the interaction potential, the relation between the Dobrushin uniqueness condition (high-temperature) and the Gaussian concentration inequality has been obtained in [11, 5, 4], whereas at low temperature weaker concentration inequalities are proved in [5].

In this paper we are interested in the behaviour of concentration inequalities under stochastic time-evolution. To our knowledge this natural question has not been addressed anywhere in the literature. There are however several motivations to be interested in this rather natural problem. First, in the context of non-equilibrium systems, non-equilibrium stationary states, or transient non-equilibrium states are usually characterized rather implicitly via an underlying dynamics. If we are interested in concentration properties of such measures, we are naturally lead to the question of time-evolution of measures satisfying a concentration inequality. It is also used in various contexts that a Markovian semigroup interpolates between different measures [1], [12, Section 2.3], and therefore it is of interest whether this interpolation conserves concentration properties. Notice that in the context of Gibbs measures, stochastic time-evolution (even high-temperature dynamics) can destroy the Gibbs property [7], therefore it is interesting to understand whether such measures – though not Gibbs – still enjoy concentration properties, or whether there can be phase transitions in the concentration behavior of a measure, *e.g.*, from Gaussian concentration bound to weaker concentration bound in a dynamics leading from high to low-temperature regime.

In this paper we focus on the so-called Gaussian concentration bound, abbreviated GCB, (see Definition 2.1 below for a precise statement), and ask under which conditions GCB is conserved under stochastic time evolution. Because we need to estimate exponential moments of a time-evolved probability measure, as we will see later on in more detail, an object popping up naturally is the so-called nonlinear semigroup $V_t(f) = \log S_t(e^f)$ where S_t is the Markov semigroup of the process under consideration, as well as its associated nonlinear generator $\mathcal{H}(f) = e^{-f} \mathcal{L}(e^f)$ where \mathcal{L} is the Markov generator. It is crucial to obtain estimates for the time-dependent Lipschitz constant of $V_t f$, which, because we can restrict to smooth f boils down to gradient estimates.

In this paper, for the stochastic dynamics, we mostly restrict to Markovian diffusion processes (only in the last section we consider non-Markovian diffusions of a specific type). In this setting, the nonlinear generator \mathcal{H} is a sum of a linear and a quadratic part, where the quadratic part coincides with the “carré du champ” operator. This implies that in the reversible setting, one can use general results on strong gradient bounds from [1], whereas in

the non-reversible setting we rely on coupling or on direct estimation of the exponential of the square distance function.

The rest of our paper is organized as follows. In section 2 we define the basic setting and define the problem of time-evolution of the Gaussian concentration bound. We also give a simple but enlightening and guiding example of the Ornstein-Uhlenbeck process, where starting from a normal distribution, we can explicitly compute the time-evolution of the constant in the Gaussian concentration bound. In section 3 we use the method of the non-linear semigroup, which as we explain in section 3.2, enters naturally in our context. The main problem is then to understand the evolution of the Lipschitz constant under the non-linear semigroup. In section 3, we control this via the method and framework of [1], using the strong gradient bound. This method applies in the reversible context. In section 5, we use a different approach based on coupling which can also be used in the non-reversible context. We give examples from non-equilibrium steady states, and non-gradient perturbations of reversible diffusions. In section 4, we use a third approach based on the exponential moment of the square distance function. With this technique, we give a class of examples where, starting from any initial measure, we have the Gaussian concentration bound at any positive time, and we also apply the technique for a time-dependent Markovian diffusion with confining drift condition. This applies for instance to the “noisy” Lorenz system. Finally, in section 6 we treat non-Markovian diffusions with linear drift, which can be studied using martingale moment inequalities. In the appendices we give a new proof of Gaussian concentration from the existence of an exponential moment of the square distance function, and provide a general approximation lemma, showing that in the context of a separable Banach space, the Gaussian concentration bound for smooth functions with bounded support implies the Gaussian concentration bound for general Lipschitz functions.

2 Setting and basic questions

2.1 Gaussian concentration bounds

We denote by $\mathcal{C}_b(\mathbb{R}^d, \mathbb{R})$ the space of bounded continuous functions from \mathbb{R}^d to \mathbb{R} . For a probability measure μ on (the Borel σ -field of) \mathbb{R}^d and $f \in \mathcal{C}_b(\mathbb{R}^d, \mathbb{R})$, we denote by $\mu(f) = \int f d\mu$ the expectation of f with respect to μ . $\text{Lip}(\mathbb{R}^d, \mathbb{R})$ denotes the set of real-valued Lipschitz functions. We

further denote for $f \in \text{Lip}(\mathbb{R}^d, \mathbb{R})$

$$\text{lip}(f) := \sup_{\substack{x, y \\ x \neq y}} \frac{|f(x) - f(y)|}{\|x - y\|}$$

the Lipschitz constant of f , where $\|\cdot\|$ denotes the Euclidean norm in \mathbb{R}^d . A Lipschitz function is almost surely differentiable by Rademacher's theorem [13, p. 101], and the supremum norm of the gradient coincides with the Lipschitz constant. For $f : \mathbb{R}^d \rightarrow \mathbb{R}$ we denote by ∇f the gradient of f , which we view as a column vector. We denote

$$\|\nabla f\|_\infty^2 := \text{ess sup}_{x \in \mathbb{R}^d} \|\nabla f(x)\|^2.$$

We can now define the notion of Gaussian concentration bound.

DEFINITION 2.1. *Let μ be a probability measure on (the Borel σ -field of) \mathbb{R}^d .*

- a) *We say that μ satisfies the smooth Gaussian concentration bound with constant D if we have*

$$\log \mu(e^{f - \mu(f)}) \leq D \text{lip}(f)^2$$

for all smooth compactly supported f . We abbreviate this property by GCBS(D).

- b) *We say that μ satisfies the Gaussian concentration bound with constant D if we have*

$$\log \mu(e^{f - \mu(f)}) \leq D \text{lip}(f)^2$$

for all Lipschitz functions $f \in \text{Lip}(\mathbb{R}^d, \mathbb{R})$. We abbreviate this property by GCB(D).

In appendix B we prove in a much more general setting, *i.e.*, in the context of a separable Banach space, that GCBS and GCB are equivalent. More precisely we prove that GCBS(D) implies GCB(D), hence the constant D does not change. (In general, we have to replace compact support by bounded support.) Therefore, for the rest of the paper, we concentrate on the time evolution of GCBS rather than GCB.

2.2 Time evolved Gaussian concentration bound

Let $\{X_t, t \geq 0\}$ denote a Markov diffusion process on \mathbb{R}^d , *i.e.*, a process solving a SDE of the form

$$dX_t = b(t, X_t) dt + \sqrt{2a(t, X_t)} dW_t. \quad (1)$$

In this equation, $b : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow \mathbb{R}^d$, $a : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow M_d^+$ (where M_d^+ denotes the set of $d \times d$ symmetric positive definite matrices), and where $\{W_t, t \geq 0\}$ is a standard Brownian motion on \mathbb{R}^d . We will also assume that b and a are as regular as needed.

The questions studied in this paper are the following.

1. If μ_0 satisfies GCBS(D_0), does the distribution μ_s at time $s > 0$ of the process $\{X_t, t \geq 0\}$, starting according to μ_0 , satisfy GCBS(D_s) for some D_s ?
2. Does the stationary measure (or stationary measures) of $\{X_t : t \geq 0\}$ satisfy GCBS(D) for some constant D ? Can one estimate D ?

2.3 Illustrative example: an Ornstein-Uhlenbeck process

A simple but inspiring example is given by the one-dimensional Ornstein-Uhlenbeck process, *i.e.*, the process $\{X_t, t \geq 0\}$ solving the SDE

$$dX_t = -\kappa X_t dt + \sigma dW_t \quad (2)$$

where $\sigma > 0$, and $\{W_t, t \geq 0\}$ is a standard Brownian motion. Let us denote by X_t^x the solution starting from $X_0 = x$. Then we have

$$X_t^x = e^{-\kappa t} x + \sigma \int_0^t e^{-\kappa(t-s)} dW_s.$$

If we start from X_0 which is normally distributed with expectation zero and variance θ^2 (denote by $\mathcal{N}(0, \theta^2)$ the corresponding distribution) then, at time $t > 0$, X_t is normally distributed with expectation zero and variance

$$\sigma_t^2 = \theta^2 e^{-2\kappa t} + \frac{\sigma^2}{2\kappa} (1 - e^{-2\kappa t}).$$

Because the normal distribution $\mathcal{N}(0, a^2)$ satisfies GCBS(D) with $D = a^2/2$ we conclude that for this example, with $\mu_0 = \mathcal{N}(0, \theta^2)$, μ_t satisfies GCBS(D_t) with

$$D_t = D_\infty + (D_0 - D_\infty) e^{-2\kappa t}$$

with $D_\infty = \frac{\sigma^2}{2\kappa}$. Hence, μ_t satisfies GCBS(D_t) with a constant D_t interpolating smoothly between the initial constant D_0 and the constant D_∞ associated to the stationary normal distribution.

In case $\kappa = 0$ the process is σB_t , and we find

$$\sigma_t^2 = \theta^2 + \sigma^2 t$$

which implies that the constant of the Gaussian concentration bound evolves as

$$D_t = D_0 + \sigma^2 t.$$

3 Nonlinear semigroup approach

In this section we develop an abstract approach based on the so-called nonlinear semigroup, combined with the Bakry-Emery Γ_2 criterion. We show that if the strong gradient bound is satisfied, then the Gaussian concentration bound is conserved in the course of the time evolution, and in the limit $t \rightarrow \infty$.

3.1 The nonlinear semigroup

Let $\{X_t : t \geq 0\}$ be a Markov diffusion process on \mathbb{R}^d as defined in (1) and denote by S_t its semigroup acting on $\mathcal{C}_b(\mathbb{R}^d, \mathbb{R})$. As usual, the generator is denoted by

$$\mathcal{L}f(x) = \lim_{t \downarrow 0} \frac{S_t f(x) - f(x)}{t}$$

on its domain $\mathcal{D}(\mathcal{L})$ of functions f such that $\frac{S_t f(x) - f(x)}{t}$ converges uniformly in x when $t \downarrow 0$. The *non-linear* semigroup is denoted by

$$V_t(f) = \log S_t(e^f).$$

This is indeed a semigroup since

$$V_{t+s}(f) = \log(S_{t+s}(e^f)) = \log(S_t(S_s(e^f))) = \log S_t(\log e^{V_s(f)}) = V_t(V_s(f)).$$

We denote by \mathcal{H} its generator, *i.e.*, for all $x \in \mathbb{R}^d$,

$$\mathcal{H}(f)(x) = \lim_{t \downarrow 0} \frac{V_t(f)(x) - f(x)}{t} \tag{3}$$

defined on the domain $\mathcal{D}(\mathcal{H})$ where the defining limit in (3) converges uniformly. The relation between \mathcal{H} and V_t is more subtle than the relation between \mathcal{L} and S_t . We will restrict ourselves to the case of diffusions with regular coefficients on \mathbb{R}^d , although what follows can be formulated in a more abstract setting. Thanks to the approximation results found in Appendix B, it is enough to restrict ourselves to adequate subsets of the domains $\mathcal{D}(\mathcal{L})$ and $\mathcal{D}(\mathcal{H})$. Denote by $\mathcal{C}_c^\infty(\mathbb{R}^d, \mathbb{R})$ the space of infinitely differentiable real-valued functions on \mathbb{R}^d with compact support.

PROPOSITION 3.1. *The following properties hold:*

1. $\mathcal{C}_c^\infty(\mathbb{R}^d, \mathbb{R}) \subset \mathcal{D}(\mathcal{L})$;
2. $\mathcal{C}_c^\infty(\mathbb{R}^d, \mathbb{R}) \subset \mathcal{D}(\mathcal{H})$, and for $f \in \mathcal{C}_c^\infty(\mathbb{R}^d, \mathbb{R})$ we have

$$\mathcal{H}(f) = e^{-f} \mathcal{L} e^f;$$

3. $\forall f \in \mathcal{C}_c^\infty(\mathbb{R}^d, \mathbb{R})$, $V_t(f) \in \mathcal{D}(\mathcal{L})$ for each $t \geq 0$;
4. $\forall f \in \mathcal{C}_c^\infty(\mathbb{R}^d, \mathbb{R})$, $V_t(f) \in \mathcal{D}(\mathcal{H})$ for each $t \geq 0$;
5. We have

$$\frac{dV_t(f)}{dt} = \mathcal{H}(V_t(f)).$$

PROOF. Property 1 is well-known, see for instance [1]. In order to prove the second property, we first observe that $\exp(-\|f\|_\infty) \leq S_t(\exp(f)) \leq \exp(\|f\|_\infty)$ and $\exp(f) \in \mathcal{D}(\mathcal{L})$ (see again [1]). Now the property follows from the definition of \mathcal{H} . To prove property 3, observe that, for each $t \geq 0$, $|V_t(f)(x)|$ is bounded in x and goes to 0 as $\|x\|$ goes to infinity. Moreover, by the usual regularity bounds, the function $x \mapsto V_t(f)(x)$ is, for each $t \geq 0$, (at least) twice differentiable with bounded derivatives (see [1]). The last two properties follow from the semigroup property of $\{V_t, t \geq 0\}$ and the fact that $S_t(\exp(f)) \in \mathcal{D}(\mathcal{L})$ for each $t \geq 0$. \square

Notice that, unlike in the case of the linear semigroup S_t , we *do not have commutation* of the semigroup with the generator, *i.e.*, in general $\mathcal{H}(V_t(f)) \neq V_t(\mathcal{H}(f))$.

3.2 Some preparatory computations

In order to start answering the questions of Section 2.2 we show here how the non-linear semigroup enters naturally into these questions. Indeed, for all $t \geq 0$, we have

$$\begin{aligned} \mu_t(e^{f-\mu_t(f)}) &= \mu_0(S_t(e^f)) e^{-\mu_0(S_t(f))} \\ &= \mu_0(e^{V_t(f)-\mu(V_t(f))}) e^{\mu_0(V_t(f)-S_t(f))}. \end{aligned} \quad (4)$$

Therefore, if μ_0 satisfies GCBS(D_0), then we can estimate the first factor in the r.h.s. of (4)

$$\mu(e^{V_t(f)-\mu(V_t(f))}) \leq e^{D_0 \text{lip}(V_t(f))^2} \quad (5)$$

and so we have to estimate $\text{lip}(V_t(f))$, which in the case of diffusion processes will boil down to estimating $\nabla V_t(f)$. Concerning the second factor in (4) we define first the “truly non-linear” part of the non-linear generator as follows

$$\mathcal{H}_{\text{nl}}(f) = \mathcal{H}(f) - \mathcal{L}(f)$$

for $f \in \mathcal{D}(\mathcal{L}) \cap \mathcal{D}(\mathcal{H})$. In the case of diffusion processes, this operator exactly contains the quadratic term of \mathcal{H} , which coincides in turn with the so-called “carré du champ operator” (see section 3.3 below).

PROPOSITION 3.2. *For regular diffusions on \mathbb{R}^d , for any $f \in \mathcal{C}_c^\infty(\mathbb{R}^d, \mathbb{R})$, for any probability measure μ_0 on \mathbb{R}^d , for all $t \geq 0$, we have*

$$\mu(|V_t(f) - S_t(f)|) \leq \|V_t(f) - S_t(f)\|_\infty \leq \int_0^t \|\mathcal{H}_{\text{nl}}(V_s(f))\|_\infty ds. \quad (6)$$

PROOF. It follows from Proposition 3.1 that

$$\begin{aligned} \frac{d(V_t(f) - S_t(f))}{dt} &= \mathcal{H}(V_t(f)) - \mathcal{L}S_t(f) = \mathcal{H}(V_t(f)) - \mathcal{L}V_t(f) + \mathcal{L}(V_t(f) - S_t(f)) \\ &= \mathcal{H}_{\text{nl}}(V_t(f)) + \mathcal{L}(V_t(f) - S_t(f)). \end{aligned}$$

As a consequence, we obtain, by the variation of constant method,

$$V_t(f) - S_t(f) = \int_0^t S_{t-s}(\mathcal{H}_{\text{nl}}(V_s(f))) ds$$

and because $\{S_t, t \geq 0\}$ is a Markov semigroup, it is a contraction semigroup in the supremum norm and because μ is a probability measure, we obtain the desired inequality. \square

As a consequence of (5) and (6), we first aim at obtaining estimates for $\text{lip}(V_t(f))$, or $\nabla V_t(f)$, and next use these estimates to further estimate the integral in the r.h.s. of (6). In particular, in the case of diffusion processes on \mathbb{R}^d , $\mathcal{H}_{\text{nl}}(g)$ is bounded in terms of $(\nabla g)^2$, and hence if we have a uniform estimate for $\nabla(V_t(f))$, we can plug it in immediately. Summarizing, assuming that μ satisfies GCBS(D), when we combine (4), (5) and (6), we obtain, for all $t \geq 0$,

$$\mu_t(e^{f - \mu_t(f)}) \leq \exp\left(D \text{lip}(V_t(f))^2 + \int_0^t \|\mathcal{H}_{\text{nl}}(V_s(f))\|_\infty ds\right). \quad (7)$$

3.3 Abstract gradient bound approach

In this subsection we study the questions formulated in Section 2.2 in the context of Markovian diffusion triples, in the sense of [1], *i.e.*, reversible diffusion processes for which we have the integration by parts formula relating the Dirichlet form and the carré du champ bilinear form. Let $\{X_t, t \geq 0\}$ be a Markov diffusion, *i.e.*, a solution of the SDE of the form (1). Moreover, we will assume in this subsection that the covariance matrix $a(x)$ is not degenerate, and is bounded, uniformly in $x, v \in \mathbb{R}^d$, *i.e.*, for some $C_1, C_2 > 0$,

$$C_1^{-2} \|v\|^2 \leq \langle v, a(x)v \rangle \leq C_2^2 \|v\|^2 \quad (8)$$

where $\langle \cdot, \cdot \rangle$ denotes Euclidean inner product.

The generator of the process $\{X_t, t \geq 0\}$ solving the SDE (1), acting on a smooth compactly supported functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is then given by

$$\mathcal{L}f(x) = \sum_{i=1}^d b_i(x) \partial_i f(x) + \sum_{i,j} a_{ij}(x) \partial_i \partial_j f(x) \quad (9)$$

where ∂_i denotes partial derivative w.r.t. x_i . We assume that the $a_{i,j}$'s and the b_i 's are regular.

To the generator \mathcal{L} is associated the carré du champ bilinear form

$$\Gamma(f, g) = \frac{1}{2} (\mathcal{L}(fg) - g\mathcal{L}(f) - f\mathcal{L}(g)) = \langle \nabla f, a \cdot \nabla g \rangle.$$

Notice that Γ satisfies the so-called diffusive condition, *i.e.*, for all smooth functions $\psi : \mathbb{R} \rightarrow \mathbb{R}$ and $x \in \mathbb{R}^d$,

$$\Gamma(\psi(f), \psi(f))(x) = (\psi')^2(f(x)) \Gamma(f, f)(x).$$

We will further assume that there exists a reversible measure ν such that the integration by parts formula

$$\int f(-\mathcal{L}g) d\nu = \int \Gamma(f, g) d\nu$$

holds. The triple $(\mathbb{R}^d, \Gamma, \nu)$ is then a Markov diffusion triple in the sense of [1, section 3.1.7].

The second order carré du champ bilinear form is given by

$$\Gamma_2(f, g) = \frac{1}{2} (\mathcal{L}\Gamma(f, f) - \Gamma(\mathcal{L}f, g) - \Gamma(f, \mathcal{L}g)).$$

In what follows, we abbreviate, as usual, $\Gamma(f, f) =: \Gamma(f)$, $\Gamma_2(f, f) = \Gamma_2(f)$. An important example is when $b = -\nabla W$ and $a = I_d$, in which case the second order the carré du champ bilinear form is given by

$$\Gamma_2(f, f) = \|\nabla \nabla f\|^2 + \langle \nabla f, \nabla \nabla W(\nabla f) \rangle$$

where $\nabla \nabla W$ denotes the Hessian of W , *i.e.*, the matrix of the second derivatives. By the non-degeneracy and boundedness condition (8), we have, for all $x \in \mathbb{R}^d$

$$C_1^{-2} \|\nabla f(x)\|^2 \leq \Gamma(f)(x) \leq C_2^2 \|\nabla f(x)\|^2.$$

Following [1] we say that the strong gradient bound is satisfied with constant $\rho \in \mathbb{R}$ if for all $t \in \mathbb{R}_+$

$$\sqrt{\Gamma(S_t f)} \leq e^{-\rho t} S_t(\sqrt{\Gamma(f)}). \quad (10)$$

This condition is fulfilled when, *e.g.*, the Bakry-Emery curvature bound,

$$\Gamma_2(f) \geq \rho \Gamma(f)$$

is satisfied. We refer to [1, Chapter 3] for the proof and more background on this formalism. We then have the following general result.

THEOREM 3.1. *Let $\{X_t, t \geq 0\}$ be a reversible diffusion process such that (10) is fulfilled. Assume that μ_0 satisfies GCBS(D_0). Then, for every $t \geq 0$, μ_t satisfies GCBS(D_t) with*

$$D_t = D_0 C_1^2 C_2^2 e^{-2\rho t} + \frac{C_1^2 C_2^4}{2\rho} (1 - e^{-2\rho t}). \quad (11)$$

In particular, if $\rho > 0$, then the unique reversible measure ν satisfies GCBS(D_∞) with $D_\infty = \frac{C_2^4 C_1^2}{2\rho}$.

PROOF. Using (10) we start by estimating $\|\nabla V_t f\|$ for $f : \mathbb{R}^d \rightarrow \mathbb{R}$ smooth with compact support

$$\begin{aligned} \|\nabla V_t(f)\| &= \frac{\|\nabla(S_t(e^f))\|}{S_t(e^f)} \leq C_1 \frac{\sqrt{\Gamma(S_t(e^f))}}{S_t(e^f)} \\ &\leq C_1 e^{-\rho t} \frac{S_t(\sqrt{\Gamma(e^f)})}{S_t(e^f)} = C_1 e^{-\rho t} \frac{S_t(e^f \sqrt{\Gamma(f)})}{S_t(e^f)} \\ &\leq C_1 e^{-\rho t} \|\sqrt{\Gamma(f)}\|_\infty \leq C_1 C_2 e^{-\rho t} \|\nabla f\|_\infty. \end{aligned}$$

As a consequence we obtain

$$\text{lip}(V_t(f)) = \|\nabla V_t f\|_\infty \leq C_1 C_2 e^{-\rho t} \|\nabla f\|_\infty. \quad (12)$$

Now we recall that what we called the “truly non-linear part” of the non-linear generator \mathcal{H}_{nl} coincides here with the carré du champ bilinear form, *i.e.*,

$$\mathcal{H}_{\text{nl}}(f) = \Gamma(f) \leq C_2^2 \|\nabla f\|_\infty^2. \quad (13)$$

As a consequence, starting from (6), we further estimate

$$\|V_t(f) - S_t(f)\|_\infty \leq C_2^2 \int_0^t \|\nabla V_s(f)\|_\infty^2 ds \leq C_1^2 C_4^2 \|\nabla f\|_\infty^2 \int_0^t e^{-2\rho s} ds. \quad (14)$$

Combining (12), (14) with (7) we obtain that μ_t satisfies GCBS(D_t) with

$$D_t = D_0 C_1^2 C_2^2 e^{-2\rho t} + C_1^2 C_2^4 \int_0^t e^{-2\rho s} ds$$

which is the claim of the theorem. \square

REMARK 3.1.

a) In case $\Gamma(f) = a^2 \|\nabla f\|^2$, we have $C_1 = a^{-2}$, $C_2 = a^2$, so D_t in $t = 0$ equals D . In general, $C_1^2 C_2^2 > 1$, which means that at time $t = 0$ we do not recover the constant D in (11), but a larger constant. This is an artefact of the method where we estimate the norm of the gradient via the carré du champ.

b) In case we have an exact commutation relation of the type

$$\nabla S_t(f) = e^{-\rho t} S_t \nabla f$$

such as is the case for the Ornstein-Uhlenbeck process, we obtain directly

$$\|\nabla V_t(f)\| \leq e^{-\rho t} \|\nabla f\|_\infty$$

i.e., without using the bilinear form Γ .

4 Coupling approach

4.1 Coupling and the nonlinear semigroup

In the previous section, the essential input coming from the strong gradient bound is the estimate (12) which implies that for all $x, y \in \mathbb{R}^d$ and all $t \in \mathbb{R}_+$

$$\|V_t(f)(x) - V_t(f)(y)\| \leq C_t \|\nabla f\|_\infty \|x - y\| e^{-\rho t}. \quad (15)$$

Once we have the bound (15), we can use it to further estimate the r.h.s. of (6), provided we have a control on \mathcal{H}_{nl} . Instead of starting from the curvature bound, in this subsection we start from a coupling point of view. This has the advantage that reversibility is no longer necessary, and moreover we can include degenerate diffusions such as the Ginzburg-Landau diffusions (see below). We denote by X_t^x the process $\{X_t, t \geq 0\}$ started at $X_0 = x$.

As an important example to keep in mind, consider the Ornstein-Uhlenbeck process on \mathbb{R}^d , with generator

$$-\langle Ax, \nabla \rangle + \Delta$$

where Δ denotes the Laplacian in \mathbb{R}^d , and where A is a $d \times d$ matrix. In that case we have

$$X_t^x = e^{-At} x + \int_0^t e^{-2A(t-s)} dW_s \quad (16)$$

which depends deterministically, and in fact linearly, on x .

DEFINITION 4.1. *Let $\gamma : [0, \infty) \rightarrow [0, \infty)$ be a measurable function such that $\gamma(0) = 1$. We say that the process $\{X_t, t \geq 0\}$ can be coupled at rate γ if, for all $x, y \in \mathbb{R}^d$, there exists a coupling of $\{X_t^x, t \geq 0\}$ and $\{X_t^y, t \geq 0\}$ such that almost surely in this coupling*

$$d(X_t^x, X_t^y) \leq d(x, y) \gamma(t). \quad (17)$$

In the case of the Ornstein-Uhlenbeck process in \mathbb{R}^d , we have from (16) (which implicitly defines a coupling, because we use (16) for all x with the same Brownian realization)

$$\|X_t^x - X_t^y\| \leq \|e^{-At}\| \|x - y\|$$

hence $\gamma(t) = \|e^{-At}\|$. Notice that $\gamma(t)$ can be “expanding” or “contracting”, depending on the spectrum of A . More precisely, γ will be eventually contracting if the numerical range of A lies in the half plane of complex numbers with non-positive real part.

REMARK 4.1. *In the context of Brownian motion on a Riemannian manifold, it is proved in [14] that (17) for $\gamma(t) = e^{-Kt/2}$ is equivalent with having K as a lower bound for the Ricci curvature, which in that context is equivalent with the Bakry-Emery curvature bound. In general however, the relation between the coupling condition (17) and the Bakry-Emery curvature bound is not so simple. In particular, the coupling approach applies beyond reversibility, in the context of degenerate diffusions, and beyond the setting of exponential decay of $\gamma(t)$ in (17).*

We have the following result. Let \mathcal{W}_1 be the space of probability measures μ such that $\int d(0, x) d\mu(x) < \infty$ equipped with the distance

$$\begin{aligned} d_{\mathcal{W}_1}(\mu, \nu) &= \sup \left\{ \int f d\mu - \int f d\nu : \text{lip}(f) \leq 1 \right\} \\ &= \inf \left\{ \int d(x, y) dP : P \text{ coupling of } \mu, \nu \right\}. \end{aligned}$$

THEOREM 4.1. *Assume that $\{X_t, t \geq 0\}$ can be coupled at rate γ . Assume that μ_0 satisfies $\text{GCBS}(D_0)$, then for all $t > 0$, and for all f smooth we have the estimate*

$$\log \mu_t(e^{f - \mu_t(f)}) \leq D_0 \text{lip}(f)^2 \gamma(t)^2 + C_2^2 \text{lip}(f)^2 \int_0^t \gamma(s)^2 ds \quad (18)$$

where C_2 is defined in (8). As a consequence, μ_t satisfies $\text{GCBS}(D_t)$ with

$$D_t = D\gamma(t)^2 + C_2^2 \int_0^t \gamma(s)^2 ds. \quad (19)$$

In particular, if $\int_0^\infty \gamma(s)^2 ds < \infty$, then every weak limit point of $\{\mu_t, t \geq 0\}$ satisfies $\text{GCBS}(D_\infty)$ with

$$D_\infty = C_2^2 \int_0^\infty \gamma(s)^2 ds.$$

Moreover, the unique invariant probability measure $\nu \in \mathcal{W}_1$ satisfies $\text{GCBS}(D_\infty)$.

PROOF. We start with a lemma which gives a general estimate on the variation of $V_t f$.

LEMMA 4.1. *Let f be Lipschitz and assume that $\{X_t, t \geq 0\}$ can be coupled at rate γ . Then for all $t \geq 0$ and $x, y \in \mathbb{R}^d$ we have*

$$V_t(f)(x) - V_t(f)(y) \leq \text{lip}(f) \gamma(t) d(x, y).$$

As a consequence, for all $t \geq 0$,

$$\text{lip}(V_t(f)) \leq \text{lip}(f) \gamma(t).$$

PROOF. Let us denote by $\widehat{\mathbb{E}}$ expectation in the coupling of $\{X_t^x, t \geq 0\}$ and $\{X_t^y, t \geq 0\}$ for which (17) holds (which exists by assumption). Then we

have

$$\begin{aligned}
\exp(V_t(f)(x) - V_t(f)(y)) &= \frac{\widehat{\mathbb{E}}(e^{f(X_t^x)})}{\widehat{\mathbb{E}}(e^{f(X_t^y)})} = \frac{\widehat{\mathbb{E}}(e^{f(X_t^y)}(e^{f(X_t^x)-f(X_t^y)}))}{\widehat{\mathbb{E}}(e^{f(X_t^y)})} \\
&\leq \frac{\widehat{\mathbb{E}}(e^{f(X_t^y)} e^{\text{lip}(f) d(X_t^x, X_t^y)})}{\widehat{\mathbb{E}}(e^{f(X_t^y)})} \\
&\leq \frac{\widehat{\mathbb{E}}(e^{f(X_t^y)} e^{\text{lip}(f) d(x,y)\gamma(t)})}{\widehat{\mathbb{E}}(e^{f(X_t^y)})} = e^{\text{lip}(f)d(x,y)\gamma(t)}
\end{aligned}$$

where in the last inequality we used (17). \square

Notice that in lemma 4.1 it is not required that $\gamma(t) \rightarrow 0$ as $t \rightarrow \infty$, *i.e.*, the coupling does not have to be successful. However if one wants to pass to the limit $t \rightarrow \infty$ then it is important that $\gamma(t) \rightarrow 0$ as $t \rightarrow \infty$. This in turn implies, as we see in the next lemma that among all probability measures in the Wasserstein space \mathcal{W}_1 , there is a unique invariant probability measure ν , and for all $\mu_0 \in \mathcal{W}_1$, $\mu_t \rightarrow \nu$ weakly as $t \rightarrow \infty$.

LEMMA 4.2. *Assume that $\{X_t, t \geq 0\}$ can be coupled at rate γ and $\gamma(t) \rightarrow 0$ as $t \rightarrow \infty$. Then there exists a unique invariant probability measure ν in \mathcal{W}_1 . Moreover, for all $\mu_0 \in \mathcal{W}_1$, $\mu_t \rightarrow \nu$ as $t \rightarrow \infty$.*

PROOF. Let μ_0, ν_0 be elements of \mathcal{W}_1 and let f be a Lipschitz function with $\text{lip}(f) \leq 1$. Because μ_0, ν_0 are elements of \mathcal{W}_1 , there exists a coupling \mathbb{P} such that

$$\int d(x, y) d\mathbb{P} = d_{\mathcal{W}_1}(\mu_0, \nu_0) < \infty.$$

Then

$$\begin{aligned}
\int f d\mu_t - \int f d\nu_t &= \int \widehat{\mathbb{E}}(f(X_t^x) - f(X_t^y)) d\mathbb{P}(x, y) \\
&\leq \int \widehat{\mathbb{E}}(d(X_t^x, X_t^y)) d\mathbb{P}(x, y) \\
&\leq \gamma(t) d_{\mathcal{W}_1}(\mu_0, \nu_0).
\end{aligned}$$

This shows that for all $\mu_0, \nu_0 \in \mathcal{W}_1$, and for all $t \geq 0$,

$$d_{\mathcal{W}_1}(\mu_t, \nu_t) \leq \gamma(t) d_{\mathcal{W}_1}(\mu_0, \nu_0). \quad (20)$$

Existence of an invariant measure $\nu \in \mathcal{W}_1$ now follows via a standard contraction argument. If $\mu, \nu \in \mathcal{W}_1$ are both invariant then (20) gives, after

taking $t \rightarrow \infty$: $d_{\mathcal{W}_1}(\mu, \nu) = 0$, which shows uniqueness of the invariant measure $\nu \in \mathcal{W}_1$. The fact that $\mu_0 \in \mathcal{W}_1$, implies $\mu_t \rightarrow \nu$ as $t \rightarrow \infty$ then also follows from (20). \square

To finish the proof of the theorem, we use (13)

$$\mathcal{H}_{\text{nl}}(f) = \Gamma(f) \leq C_2^2 \|\nabla f\|^2 \leq C_2^2 \text{lip}(f)^2.$$

Combining with (6) and (5) and lemma 4.2 this yields the result of the theorem.

\square

As an application we have the following result on Markovian diffusions with covariance matrix a not depending on the location x .

THEOREM 4.2. *Let X_t denote a diffusion process on \mathbb{R}^d with generator of type (9), and where the covariance matrix a does not depend on location x . Assume furthermore that the function $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is continuously differentiable and the differential $D_x b$ satisfies the estimate*

$$\langle D_x b(x)(u), u \rangle \leq -\kappa \|u\|^2 \quad (21)$$

for all $x, u \in \mathbb{R}^d$ and some $\kappa \in \mathbb{R}$. Let μ_0 satisfy $\text{GCBS}(D_0)$, then, for all $t > 0$, μ_t satisfies $\text{GCBS}(D_t)$ with

$$D_t = D_0 e^{-2\kappa t} + \frac{\|a\|}{2\kappa} (1 - e^{-2\kappa t}). \quad (22)$$

Moreover, if $\kappa > 0$, then $\mu_t \rightarrow \nu$ as $t \rightarrow \infty$ where ν is the unique invariant probability measure, which satisfies $\text{GCB}(\|a\|/2\kappa)$. In particular, if $b = -\nabla W$, where the potential $W : \mathbb{R}^d \rightarrow \mathbb{R}$ is \mathcal{C}^2 , then (21) reduces to the convexity condition

$$\langle \nabla \nabla W, u, u \rangle \geq \kappa \|u\|^2.$$

PROOF. We have $\|\mathcal{H}_{\text{nl}}(f)\| = \Gamma(f) \leq \|a\|(\nabla f)^2$. Therefore by Theorem 4.1 it suffices to see that we have a coupling rate $\gamma(t) = e^{-\kappa t}$. We couple X_t^x, X_t^y by using the same realization of the underlying Brownian motion $\{W_t, t \geq 0\}$, and as a consequence, because a does not depend on x , the difference $X_t^x - X_t^y$ is evolving according to

$$\frac{d(X_t^x - X_t^y)}{dt} = b(X_t^x) - b(X_t^y).$$

We have

$$\begin{aligned} b(X_t^x) - b(X_t^y) &= \int_0^1 \frac{d}{ds} (b(sX_t^x + (1-s)X_t^y)) ds \\ &= \int_0^1 D_x b(\xi(s))(X_t^x - X_t^y) ds \end{aligned}$$

with

$$\xi(s) = sX_t^x + (1-s)X_t^y.$$

Therefore

$$\frac{d(\|X_t^x - X_t^y\|^2)}{dt} = 2 \int_0^1 \langle X_t^x - X_t^y, D_x b(\xi(s))(X_t^x - X_t^y) \rangle ds \leq -2\kappa \|X_t^x - X_t^y\|^2$$

which implies

$$\|X_t^x - X_t^y\| \leq e^{-\kappa t} \|x - y\|$$

for all $x, y \in \mathbb{R}^d$ and for all $t \in \mathbb{R}_+$. \square

REMARK 4.2.

a) Notice that in the approach based on the strong gradient bound, we needed non-degeneracy of the covariance matrix a in (1), cf. condition (8). In the coupling setting, we do allow the matrix a to be degenerate, but not depending on x , and the condition is only on the drift b .

b) Unlike the time dependent constant D_t , given via the strong gradient bound (11), the bound (22) yields the correct constant D at time zero. Remark that the constant of the limiting stationary distribution, which is $\|a\|/2\kappa$, is invariant under linear rescaling of time, as it should. More precisely, if we multiply the generator with a factor α , $\|a\|$ is multiplied by this same factor α , and so is the constant κ .

c) Note that inequality 21 for all x and y is equivalent to

$$\langle b(x) - b(y), x - y \rangle \leq -\kappa \|x - y\|^2$$

for all x, y .

4.2 Examples

Example 1: Ornstein-Uhlenbeck process and Brownian motion. Coming back to the simple example of the Ornstein-Uhlenbeck process (2), we have coupling rate

$$\gamma(t) = e^{-\kappa t}$$

and we find (18), *i.e.*, the time evolution of the constant in the Gaussian concentration bound is the same in general as for the special case of a Gaussian starting measure. If we have a standard Brownian motion, then the coupling rate $\gamma(t) = 1$ and the formula (19) reads ($\|a\| = 1$)

$$D_t = D_0 + t$$

which is sharp if the starting measure is the normal law $\mu_0 = \mathcal{N}(0, \sigma^2)$, which at time t gives $\mu_t = \mathcal{N}(0, \sigma^2 + t)$.

Example 2: Ginzburg-Landau dynamics with boundary reservoirs. We consider the process $\{X_t, t \geq 0\}$ on \mathbb{R}^N with generator

$$\mathcal{L} = \sum_{i=1}^N (\partial_i - \partial_{i+1})^2 - (x_{i+1} - x_i)(\partial_{i+1} - \partial_i) + L_1 + L_N$$

where ∂_i denotes partial derivative w.r.t. x_i , and where the extra operators L_1 and L_N model the reservoirs and are given by

$$L_1 = b_1(x_1) \partial_1 + \frac{1}{2} \sigma_1^2 \partial_1^2$$

$$L_N = b_N(x_N) \partial_N + \frac{1}{2} \sigma_N^2 \partial_N^2.$$

Here, $\sigma_1, \sigma_N > 0$, and the drifts associated to the reservoirs $b_1, b_N : \mathbb{R} \rightarrow \mathbb{R}$ are smooth functions. Such diffusion processes are studied in the literature on hydrodynamic limits, see *e.g.* [9].

This models a non-equilibrium transport system driven by reservoirs with drift b_1, b_N . In absence of the reservoir driving, the bulk system with generator $\sum_{i=1}^N (\partial_i - \partial_{i+1})^2 - (x_{i+1} - x_i)(\partial_{i+1} - \partial_i)$ has reversible measures which are products of mean zero Gaussians. If $b_1(x_1) = -\alpha x_1, b_N(x_N) = -\alpha x_N$, with $\alpha_1, \alpha_N > 0$, then the system is in equilibrium with reversible Gaussian product measure $C \exp(-\alpha \sum_{i=1}^N x_i^2)$.

In all other cases, by the coupling to the reservoirs, a non-equilibrium steady state is created.

For the choice $b_1(x) = -\alpha_1 x, b_N(x) = -\alpha_N x$, with $\alpha_1 \neq \alpha_N$ this corresponds to a “non-equilibrium” Ornstein-Uhlenbeck process, for which it can be shown that the unique stationary measure μ is a product of mean zero Gaussians, with variance given by

$$\int x_i^2 d\mu(x) = \frac{1}{\alpha_1} + \left(\frac{1}{\alpha_N} - \frac{1}{\alpha_1} \right) \frac{i}{N+1}$$

linearly interpolating between the left and right reservoirs.

The noise in the system is degenerate, but does not depend on x , which means that the coupling condition is satisfied. The covariance matrix a of (1) is given by $a_{ii} = 2, 2 \leq i \leq N - 1, a_{11} = 1, a_{NN} = 1, a_{i,i+1} = -1, 1 \leq i \leq N - 1$.

If the drifts associated to the reservoirs b_1, b_N are not linear, then the stationary non-equilibrium state is unknown and not Gaussian. In the following, direct application of Theorem 4.2 then gives the following.

PROPOSITION 4.1. *If the reservoir drifts are such that, for some $\kappa_N > 0$, and for all $u \in \mathbb{R}^N$,*

$$\langle u, -\Delta u \rangle - u_1^2 b'_1(x_1) - u_N^2 b'_N(x_N) \leq -\kappa_N \|u\|^2$$

with $-\Delta$ the discrete laplacian defined via $(\Delta u)_i = u_{i+1} + u_{i-1} - 2u_i$ for $2 < i < N - 1$, and $(\Delta u)_1 = u_2 - u_1, (\Delta u)_N = u_{N-1} - u_N$, then the unique stationary measure of the process with generator \mathcal{L} satisfies GCBS(D), with $D = C_N^2/2\kappa_N$, with $C_N = \|a\| \leq 4$.

Example 3: Perturbation of the drift. Remark that if (21) is satisfied with $\kappa > 0$ for the drift b with constant κ and \tilde{b} is such that $\langle D_x(\tilde{b} - b)(u), u \rangle \leq \epsilon \|u\|^2$, for some $0 < \epsilon < \kappa$, then obviously, (21) is satisfied for the drift \tilde{b} with constant $\tilde{\kappa} = \kappa - \epsilon$. For instance, if $\tilde{b}(x) = -\nabla W(x) + \epsilon(x)$, where $W(x)$ is a strictly convex potential, then if $\|D_x \epsilon\|_\infty$ is sufficiently small, there is a unique invariant probability measure ν which satisfies GCBS. However, ϵ is allowed to be of non-gradient form, which implies that ν is not known in explicit form. The same applies to systems where one adds sufficiently weak “boundary” reservoirs as long as the noise of these reservoirs does not depend on x .

5 Distance Gaussian moment approach

In this section, we start with a different approach, based on the equivalence between GCBS(D) and the existence of a Gaussian estimate of an exponential moment of the square of the distance (cf. Theorem 5.1 below).

5.1 A general equivalence

In this subsection, we work in a general separable metric space (Ω, d) . We first generalize Definition 2.1.

DEFINITION 5.1. Let μ be a probability measure on (the Borel σ -field of) (Ω, d) . We say that μ satisfies a Gaussian concentration bound with constant $D > 0$ on the metric space (Ω, d) if there exists $x_0 \in \Omega$ such that $\int d(x_0, x) d\mu(x) < \infty$ and for all $f \in \text{Lip}(\Omega, \mathbb{R})$, one has

$$\int e^{f - \mu(f)} d\mu \leq e^{D \text{lip}(f)^2}.$$

For brevity we shall say that μ satisfies $\text{GCB}(D)$ on (Ω, d) .

REMARK 5.1.

a) Note that if there exists $x_0 \in \Omega$ such that $\int d(x_0, x) d\mu(x) < \infty$ then, by the triangle inequality, $\int d(x_0, x) d\mu(x) < \infty$ for all $x_0 \in \Omega$, and all Lipschitz functions on (Ω, d) are μ -integrable.

b) Note that one can find a topological space and a probability on the Borel sigma-algebra and two distances d_1 and d_2 s.t. μ satisfies GCB on the metric space with d_1 but it does not on the metric space with d_2 . For example, take \mathbb{R} , μ to be the Gaussian measure, d_1 the Euclidean distance and $d_2(x, y) = \int_x^y (1 + |s|) ds$.

THEOREM 5.1. Let μ a probability measure on (Ω, d) . Then μ satisfies a Gaussian concentration bound if and only if it has a Gaussian moment. More precisely, we have the following:

1. If μ satisfies $\text{GCB}(D)$, there exists $x_0 \in \Omega$ such that

$$\int e^{\frac{d(x_0, x)^2}{16D}} d\mu(x) \leq 3 e^{\frac{\mu(d)^2}{D}} \quad (23)$$

where $\mu(d) := \int d(x, x_0) d\mu(x)$.

2. If there exist $x_0 \in \Omega$, $a > 0$ and $b \geq 1$ such that

$$\int e^{ad(x_0, x)^2} d\mu(x) \leq b \quad (24)$$

then μ satisfies $\text{GCB}(D)$ with

$$D = \frac{1}{2a} \left(1 \vee \frac{b^2 e}{2\sqrt{\pi}} \right). \quad (25)$$

This result can be found in [8, Theorem 2.3] with less explicit constants. We provide a direct proof of the theorem in appendix A. Notice that, by the triangle inequality, if (23) holds for some x_0 then it holds for any x_0 (with possibly different constants). The same result holds for (24).

5.2 Example 1: Diffusions coming down from infinity

As a first example of application, we consider diffusions “coming down from infinity” for which we show that from any starting measure, at positive times $t > 0$, $\text{GCBS}(D)$ holds. We refer to [2] for more on diffusions “coming down from infinity” in the one dimensional case.

We consider a diffusion process on \mathbb{R}^d which solves the SDE (1) and satisfies for some $C_1, C_2 > 0$ and for any $t \geq 0$, $x \in \mathbb{R}^d$ and $v \in \mathbb{R}^d$

$$C_1 \|v\|^2 \leq \langle v, a(t, x)v \rangle \leq C_2 \|v\|^2 \quad (26)$$

We introduce the following condition on the drift.

CONDITION 5.1. *There exists a real, non-negative, non-decreasing and \mathcal{C}^1 function h and a constant $A > 0$ such that for all $x \in \mathbb{R}^d$ and all $t \geq 0$*

$$\frac{\langle x, b(t, x) \rangle}{\|x\|} \leq A - h(\|x\|). \quad (27)$$

THEOREM 5.2. *Under condition 5.1, if additionally we have the integrability condition*

$$\int_0^\infty \frac{du}{h(u)} < \infty \quad (28)$$

then there exists $t_ > 0$, a non-negative function $C(t)$ and a constant $\alpha > 0$ such that for all $0 \leq t \leq t_*$*

$$\sup_{x \in \mathbb{R}^d} \mathbb{E}_x \left(e^{\alpha \|X_t\|^2} \right) \leq C(t).$$

We deduce the following result showing immediate Gaussian concentration in the course of diffusions coming down from infinity.

THEOREM 5.3. *Assume that condition 5.1 and (28) hold. Let μ_0 be any probability measure on (the Borel field of) \mathbb{R}^d . Let t_* and $C(t)$ be as in Theorem 5.2. Then, for all $t > 0$, the probability measure $(\mu_t)_{t \geq 0}$ defined by*

$$\mu_t(f) = \mathbb{E}_{\mu_0}(f(X_t)), \forall f \in \mathcal{C}_b(\mathbb{R}^d)$$

satisfies $\text{GCB}(D_t)$ where

$$D_t = \begin{cases} \frac{1}{2\alpha} \left(1 \vee \frac{C(t)e}{2\sqrt{\pi}} \right) & \text{if } 0 < t < t_* \\ \frac{1}{2\alpha} \left(1 \vee \frac{C(t_*)e}{2\sqrt{\pi}} \right) & \text{if } t \geq t_* . \end{cases}$$

PROOF. For $0 < t \leq t_*$, the result follows from Theorems 5.2 and 5.1. For $t > t_*$ the result follows recursively. Namely, assume that for any $0 < t \leq k t_*$ where $k > 0$ is an integer we have

$$\sup_{x \in \mathbb{R}^d} \mathbb{E}_x \left(e^{\alpha \|X_t\|^2} \right) \leq C(t)$$

where $C(t) = C(t_*)$ for $t \geq t_*$. For any $t \in [k t_*, (k+1) t_*]$, we can apply Theorem 5.2 to μ_{t-t_*} since $0 < t - t_* \leq k t_*$ and we extend the previous bound to the time interval $]0, (k+1) t_*]$, and hence recursively to any $t > 0$. The result follows from Theorem 5.1. \square

REMARK 5.2. *Theorem 5.3 implies tightness of the family $(\mu_t)_{t>0}$. Therefore the stochastic process (X_t) has invariant probability measures, each of them satisfying GCB(D_{t_*}). By standard arguments one can show that there is a unique invariant probability measure and it is absolutely continuous with respect to the Lebesgue measure.*

PROOF of Theorem 5.2. Define

$$u(t, x) = \varphi(t) e^{\alpha \|x\|^2}$$

where α and φ will be chosen later on. We have using condition 5.1 and (26)

$$\begin{aligned} & \partial_t u(t, x) + \mathcal{L}u(t, x) \\ &= e^{\alpha \|x\|^2} \left(\dot{\varphi}(t) + \varphi(t) [2\alpha \operatorname{Tr}(a(t, x)) + 4\alpha^2 \langle x, a(t, x)x \rangle + 2\alpha \langle x, b(t, x) \rangle] \right) \\ &\leq e^{\alpha \|x\|^2} \left(\dot{\varphi}(t) + \varphi(t) [2d\alpha C_2 + 4\alpha^2 C_2 \|x\|^2 + 2\alpha A \|x\| - 2\alpha h(\|x\|) \|x\|] \right). \end{aligned}$$

Using integration by parts we get

$$\int_0^z \frac{du}{h(u)} = \frac{z}{h(z)} + \int_0^z \frac{u h'(u)}{h(u)^2} du$$

and using that h is non-decreasing we obtain

$$\liminf_{z \rightarrow \infty} \frac{h(z)}{z} \geq \frac{1}{\int_0^\infty \frac{du}{h(u)}} > 0.$$

Therefore, choosing $1/2 > \alpha > 0$ sufficiently small and $y_* > 0$ sufficiently large, we have for $u \geq y_*$

$$2\alpha h(u) - 4\alpha^2 u C_2 - 2\alpha A - \frac{dC_2 \alpha}{u} > \alpha h(u).$$

Hence if $\|x\| > y_*$ we obtain

$$\partial_t u(t, x) + \mathcal{L}u(t, x) \leq u(t, x) \left(\dot{\varphi}(t) - \alpha \varphi(t) h(\|x\|) \|x\| \right). \quad (29)$$

We then define a non-increasing function $y(s)$ and the non-decreasing function $\varphi(s)$ via

$$\frac{\dot{\varphi}(s)}{\varphi(s)} = -\dot{y}(s) y(s) = \alpha y(s) \frac{h(y(s))}{2}.$$

Imposing additionally $y(0) = \infty$ we obtain

$$\int_{y(s)}^{\infty} \frac{du}{h(u)} = \frac{\alpha s}{2}.$$

We define t_* via

$$\int_{y_*}^{\infty} \frac{du}{h(u)} = \frac{\alpha t_*}{2}$$

and for $0 \leq s \leq t_*$

$$\varphi(s) = e^{-y(s)^2/2}.$$

Note that for $0 \leq s \leq t_*$ we have $y(s) \geq y_*$. Let $B > 2 \max\{y_*, \|x\|\}$. Using Itô's formula with T_B the hitting time of the boundary of the ball centered at x with radius B , we get

$$\mathbb{E}_x(u(t \wedge T_B, X_{t \wedge T_B})) = \mathbb{E}_x \left(\int_0^{t \wedge T_B} (\partial_t u + \mathcal{L}u)(s, X_s) ds \right).$$

For $0 \leq s \leq t_*$, if $\|X_s\| \geq y(s) \geq y_*$ we have using (29) and the monotonicity of h

$$(\partial_t u + \mathcal{L}u)(s, X_s) \leq e^{\alpha \|X_s\|^2} \left(\dot{\varphi} - \alpha \varphi \frac{h(y(s)y(s))}{2} \right) = 0.$$

For $0 \leq s \leq t_*$, if $\|X_s\| < y(s)$ we have

$$(\partial_t u + \mathcal{L}u)(s, X_s) \leq C e^{\alpha y(s)^2} (\dot{\varphi}(s) + \varphi(s) (1 + y(s)^2))$$

for some (computable) constant $C > 0$ independent of s . Therefore if $0 \leq t \leq t_*$ we obtain

$$\begin{aligned} & \mathbb{E}_x \left(\int_0^{t \wedge T_B} (\partial_t u + \mathcal{L}u)(s, X_s) ds \right) \\ & \leq C \mathbb{E}_x \left(\int_0^{t \wedge T_B} e^{\alpha y(s)^2} (\dot{\varphi}(s) + \varphi(s) (1 + y(s)^2)) ds \right) \\ & \leq C \int_0^t e^{\alpha y(s)^2} (\dot{\varphi}(s) + \varphi(s) (1 + y(s)^2)) ds \\ & = -C \int_0^t e^{\alpha y(s)^2} \dot{y}(s) y(s) e^{-y(s)^2/2} ds + C \int_0^t e^{\alpha y(s)^2} e^{-y(s)^2/2} (1 + y(s)^2) ds \end{aligned}$$

and since $\alpha < 1/2$

$$\leq C \int_{y(t)}^{\infty} e^{\alpha y^2} y e^{-y^2/2} dy + \mathcal{O}(1) C \int_0^t ds = \frac{C}{1-2\alpha} e^{-(1-2\alpha)y(t)^2/2} + \frac{2C}{1-2\alpha} t.$$

We now observe that since $u \geq 0$, for any $0 \leq t \leq t_*$ we have

$$\mathbb{E}_x(u(t, X_t) \mathbf{1}_{\{T_B > t\}}) \leq \mathbb{E}_x(u(t \wedge T_B, X_{t \wedge T_B})) \leq \frac{C}{1-2\alpha} e^{-(1-2\alpha)y(t)^2/2} + \frac{2C}{1-2\alpha} t.$$

Therefore by the monotone convergence theorem (letting B tend to infinity)

$$\mathbb{E}_x(u(t, X_t)) \leq \frac{C}{1-2\alpha} e^{-(1-2\alpha)y(t)^2/2} + \frac{2C}{1-2\alpha} t.$$

The result follows with

$$C(t) = \frac{C}{(1-2\alpha)} \left(e^{\alpha y(t)^2} + t e^{y(t)^2/2} \right).$$

□

5.3 Example 2: Markovian diffusion processes with space-time dependent drift and covariance

In this section, we consider diffusions which do not come down from infinity as strongly as in Theorem 5.2, for example the Ornstein-Uhlenbeck process. The setting is that of stochastic differential equations on \mathbb{R}^d given by

$$dX_t = b(X_t, t) dt + \sigma(X_t, t) dW_t$$

where the vector field b and the matrix-valued σ are regular in x, t . We assume that, for any given initial condition x_0 , the solution exists, is unique and defined for all times. This generalizes the coupling setting of Theorem 4.2, *i.e.*, we impose a more general confining condition on the drift $b(x, t)$ and allow the covariance matrix $\sigma(x, t)$ to depend on time and location.

THEOREM 5.4. *Assume that $\alpha > 0$, $\beta > 0$ and $\theta > 0$ such that, for all $x \in \mathbb{R}^d$ and $t \geq 0$*

$$\langle x, b(x, t) \rangle \leq \alpha \|x\| - \beta \|x\|^2 \tag{30}$$

and

$$\sigma^t(x, t) \sigma(x, t) \leq \theta I_d$$

where the second inequality is in the sense of the order on positive definite matrices. Then, for every initial probability measure μ_0 on \mathbb{R}^d satisfying

GCBS(D_0), the evolved probability measure μ_t satisfies GCB(D_t) for all $t \geq 0$, where D_t is given by the formula (25), with

$$\begin{aligned} a &= a_0 = \frac{\beta}{2\theta} \wedge \frac{1}{16D_0} \\ b &= b_t = b_0 \exp\left(-a_0 \left(\theta d + \frac{2\alpha^2}{\beta}\right) t\right) \\ &\quad + 2e^{\frac{4a_0}{\beta}(\theta d + \frac{2\alpha^2}{\beta})} \left(1 - \exp\left(-a_0 \left(\theta d + \frac{2\alpha^2}{\beta}\right) t\right)\right) \end{aligned}$$

and

$$b_0 = 3e^{\mu_0(d)^2/8D}$$

where $\mu_0(d) = \int \|x\| d\mu(x)$.

The same conclusions as in Remark 5.2 hold with GCB(D_∞) where

$$D_\infty = \frac{1}{2a} \left(1 \vee \frac{b_\infty^2 e}{2\sqrt{\pi}}\right).$$

PROOF. Let $a_0 = \frac{\beta}{2\theta} \wedge \frac{1}{16D_0}$ and define $u(x) = e^{a_0\|x\|^2}$. Using the assumptions we get

$$\begin{aligned} \mathcal{L}u(x) &\leq (2a_0^2\theta\|x\|^2 + a_0\theta d + 2a_0\alpha\|x\| - 2a_0\beta\|x\|^2)u(x) \\ &\leq a_0(\theta d + 2\alpha\|x\| - \beta\|x\|^2)u(x) \\ &\leq a_0 \left(\theta d + \frac{2\alpha^2}{\beta} - \frac{\beta}{2}\|x\|^2\right) u(x). \end{aligned}$$

For any $A > 0$, let $T_A = \inf\{t \geq 0 : \|X_t\| \geq A\}$. Using Dynkin's formula and Theorem 5.1, we get

$$\begin{aligned} &\mathbb{E}_{\mu_0} \left(e^{a_0\|X_{t \wedge T_A}\|^2} \right) \\ &\leq b_0 + a_0 \mathbb{E}_{\mu_0} \left(\int_0^{t \wedge T_A} e^{a_0\|X_s\|^2} \left(\theta d + \frac{2\alpha^2}{\beta} - \frac{\beta}{2}\|X_s\|^2 \right) ds \right) \end{aligned} \quad (31)$$

where, via (23)

$$b_0 = \int e^{a_0\|x\|^2} d\mu(x) \leq 3e^{\frac{\mu(d)^2}{8D}}$$

where $\mu(d) = \int \|x\| d\mu(x)$. We now estimate the expectation on the right-hand side of (31). Define, for $s > 0$, the event

$$\mathcal{E}_s = \left\{ \|X_s\|^2 > \frac{4}{\beta}\theta d + \frac{2\alpha^2}{\beta} \right\}.$$

We have

$$\begin{aligned}
& \mathbb{E}_{\mu_0} \left(e^{a_0 \|X_{t \wedge T_A}\|^2} \right) \\
& \leq b_0 + 2a_0 \left(\theta d + \frac{2\alpha^2}{\beta} \right) \mathbb{E}_{\mu_0} \left(\int_0^{t \wedge T_A} e^{a_0 \|X_s\|^2} \mathbb{1}_{\mathcal{E}_s^c} ds \right) \\
& \quad - a_0 \left(\theta d + \frac{2\alpha^2}{\beta} \right) \mathbb{E}_{\mu_0} \left(\int_0^{t \wedge T_A} e^{a_0 \|X_s\|^2} ds \right) \\
& \leq b_0 + 2a_0 \left(\theta d + \frac{2\alpha^2}{\beta} \right) e^{\frac{4a_0}{\beta} \left(\theta d + \frac{2\alpha^2}{\beta} \right) t} \\
& \quad - a_0 \left(\theta d + \frac{2\alpha^2}{\beta} \right) \mathbb{E}_{\mu_0} \left(\int_0^{t \wedge T_A} e^{a_0 \|X_s\|^2} ds \right).
\end{aligned}$$

By the Monotone Convergence Theorem, letting $A \uparrow \infty$, and Fubini's Theorem, we get

$$\begin{aligned}
& \mathbb{E}_{\mu_0} \left(e^{a_0 \|X_t\|^2} \right) \\
& \leq b_0 + 2a_0 \left(\theta d + \frac{2\alpha^2}{\beta} \right) e^{\frac{4a_0}{\beta} \left(\theta d + \frac{2\alpha^2}{\beta} \right) t} \\
& \quad - a_0 \left(\theta d + \frac{2\alpha^2}{\beta} \right) \int_0^t \mathbb{E}_{\mu_0} \left(e^{a_0 \|X_s\|^2} \right) ds.
\end{aligned}$$

Using Grönwall's lemma, we obtain

$$\begin{aligned}
\mathbb{E}_{\mu_0} \left(e^{a_0 \|X_t\|^2} \right) & \leq b_0 \exp \left(-a_0 \left(\theta d + \frac{2\alpha^2}{\beta} \right) t \right) + \\
& \quad 2 e^{\frac{4a_0}{\beta} \left(\theta d + \frac{2\alpha^2}{\beta} \right) t} \left(1 - \exp \left(-a_0 \left(\theta d + \frac{2\alpha^2}{\beta} \right) t \right) \right).
\end{aligned}$$

By Theorem 5.1, we deduce that μ_t satisfies GCBS(D_t) with the announced constant D_t . \square

REMARK 5.3.

In the case of diffusions coming down from infinity, we saw in Theorem 5.2 that GCB develops out of the time evolution of any initial distribution. In Theorem 5.4 we required that the initial distribution satisfies GCB. In the case of the one-dimensional Ornstein-Uhlenbeck process, if one starts for example with the initial probability distribution

$$d\mu_0(x) = \frac{\sqrt{2} dx}{\pi(1+x^4)}$$

it is easy to verify by an explicit computation that for any $t \geq 0$ and any $a > 0$

$$\int_{-\infty}^{\infty} d\mu_0(x) \mathbb{E}_x \left(e^{a X_t^2} \right) = \infty .$$

As an application, we consider the famous Lorenz system

$$\begin{cases} \frac{dx}{dt} &= \sigma(y - x) \\ \frac{dy}{dt} &= rx - y - xz \\ \frac{dz}{dt} &= xy - bz \end{cases}$$

which, for a certain range of (strictly positive) parameters has a strange attractor [10, Chapter 14].

Adding a noise which satisfies the condition of Theorem 4.2, this leads to a unique invariant probability measure whose properties are largely unknown. However, this measure satisfies GCBS. This can be proved observing that the Lorenz system translated by the vector $(0, 0, -2r)$ satisfies (30) using the squared norm $\|(x, y, z)\|^2 = rx^2 + \sigma y^2 + \sigma z^2$ with

$$\beta = \inf \frac{rx^2 + y^2 + bz^2}{rx^2 + \sigma y^2 + \sigma z^2} = \min \{1, \sigma^{-1}, b\sigma^{-1}\}$$

where the infimum is taken over x, y, z in such that $(x, y, z) \neq (0, 0, 0)$.

6 Non Markovian diffusions: Martingale moment approach

In this section we consider the simplest context beyond the Markov case, where we can no longer rely on methods based on generators.

We consider the stochastic differential equation on \mathbb{R} given by

$$dX_t = -\kappa X_t dt + \sigma_t dW_t \tag{32}$$

where we assume that the process σ_t is uniformly bounded and predictable. An example of this setting is

$$\begin{cases} dY_t &= -\theta Y_t + dW_t \\ dX_t &= -\kappa X_t + \sigma(Y_t) dW_t . \end{cases}$$

Then the couple (X_t, Y_t) is Markov but X_t is not, and satisfies a SDE of the form (32).

Because the process $\{X_t, t \geq 0\}$ is no longer a Markov process (unless σ_t depends only on X_t) we can no longer use techniques based on the generator as we did before for processes of Ornstein-Uhlenbeck type. The main point is that as a consequence, X_t^x equals a *deterministic process* of bounded variation plus a stochastic integral w.r.t. dW_t . As a consequence, the Gaussian concentration bound can be obtained from estimating the stochastic integral, which can be done with the help of Burkholder's inequalities.

The assumption (32) allows us to write the solution in the form

$$X_t = X_0 e^{-\kappa t} + \int_0^t e^{-\kappa(t-s)} \sigma_s dW_s. \quad (33)$$

We have the following result.

THEOREM 6.1. *Assume that there exists $M > 0$ such that*

$$\sup_{t \geq 0} \|\sigma_t\|_{L^\infty} \leq M.$$

Assume X_0 is distributed according to a probability measure μ_0 satisfying GCBS(D_0). Then we have that for all $t > 0$ there exists $D_t > 0$ such that X_t satisfies GCBS(D_t). Moreover, if $\kappa > 0$ then all weak limit points of $\{X_t, t \geq 0\}$ satisfy GCBS(D_∞) for some $D_\infty > 0$.

PROOF. We use Theorem 5.1, and will prove that there exist $a > 0, b > 0$ such that

$$\mathbb{E}_{\mu_0} \left(e^{aX_t^2} \right) \leq b.$$

Then we can conclude via Theorem 5.1, that the distribution of X_t satisfies GCBS(C) with $C \leq \frac{1}{2a} (1 \vee \frac{b^2 e}{2\sqrt{\pi}})$. We start from (33) from which we derive the inequality

$$X_t^2 \leq 2X_0^2 e^{-2\kappa t} + 2 \left(\int_0^t e^{-\kappa(t-s)} \sigma_s dW_s \right)^2. \quad (34)$$

We start by estimating, for $\gamma > 0$

$$\begin{aligned} & \mathbb{E}_{\mu_0} \left[\exp \left(\gamma \left(\int_0^t e^{-\kappa(t-s)} \sigma_s dW_s \right)^2 \right) \right] \\ &= \sum_{n=0}^{\infty} \frac{\gamma^n}{n!} \mathbb{E}_{\mu_0} \left[\left(\int_0^t e^{-\kappa(t-s)} \sigma_s dW_s \right)^{2n} \right]. \end{aligned}$$

Next use Burkholder's inequality [6] which states that for a martingale $\{Z_t, t \geq 0\}$ w.r.t. Brownian filtration, with quadratic variation $[Z, Z]_t$, we have the estimate

$$\mathbb{E}_{\mu_0}(Z_t^{2n}) \leq A(2n)^n \mathbb{E}_{\mu_0}([Z, Z]_t^n)$$

with A an absolute constant. As a consequence, we get

$$\begin{aligned} \mathbb{E}_{\mu_0} \left[\left(\int_0^t e^{-\kappa(t-s)} \sigma_s dW_s \right)^{2n} \right] &= e^{-2n\kappa t} \mathbb{E}_{\mu_0} \left[\left(\int_0^t e^{\kappa s} \sigma_s dW_s \right)^{2n} \right] \\ &\leq e^{-2n\kappa t} A(2n)^n \mathbb{E}_{\mu_0} \left[\left(\int_0^t e^{2\kappa s} \sigma_s^2 ds \right)^n \right] \\ &\leq e^{-2n\kappa t} AM^{2n} (2n)^n \mathbb{E}_{\mu_0} \left[\left(\int_0^t e^{2\kappa s} ds \right)^n \right] \\ &\leq AM^{2n} (2n)^n \left(\frac{1 - e^{-2\kappa t}}{2\kappa} \right)^n. \end{aligned}$$

As a consequence we obtain

$$\begin{aligned} &\mathbb{E}_{\mu_0} \left[\exp \left(\gamma \left(\int_0^t e^{-\kappa(t-s)} \sigma_s dW_s \right)^2 \right) \right] \\ &\leq A \sum_{n=0}^{\infty} \frac{1}{n!} \gamma^n M^{2n} (2n)^n \left(\frac{1 - e^{-2\kappa t}}{2\kappa} \right)^n. \end{aligned}$$

The r.h.s. of this inequality is a convergent series provided

$$\gamma < \left(2eM^2 \left(\frac{1 - e^{-2\kappa t}}{2\kappa} \right) \right)^{-1}.$$

We then estimate, using (34) and the Cauchy-Schwarz inequality

$$\mathbb{E}_{\mu_0} [e^{aX_t^2}] \leq \left(\mathbb{E}_{\mu_0} [e^{4aX_0^2} e^{-2\kappa t}] \right)^{1/2} \left(\mathbb{E}_{\mu_0} [e^{4a \left(\int_0^t e^{-\kappa(t-s)} \sigma_s dW_s \right)^2}] \right)^{1/2}. \quad (35)$$

Because by assumption the distribution of X_0 satisfies GCBS(C), we have that the first factor in the r.h.s. in (35) is finite as soon as $4a e^{-2\kappa t} < a_0$ where a_0 is such that $\mathbb{E}_{\mu_0}(e^{a_0 X_0^2}) < \infty$. The second factor is finite as soon as

$$a < \left(8eM^2 \left(\frac{1 - e^{-2\kappa t}}{2\kappa} \right) \right)^{-1}.$$

Therefore, $\mathbb{E}_{\mu_0}(e^{aX_t^2})$ is finite for

$$a < \left(8eM^2 \left(\frac{1 - e^{-2\kappa t}}{2\kappa} \right) \right)^{-1} \wedge a_0 e^{2\kappa t}$$

which, combined with Theorem 5.1, concludes the proof of the theorem. \square

A Proof of Theorem 5.1

Statement 1. Choose $x_0 \in \Omega$ arbitrarily. Since $x \mapsto d(x_0, x)$ is 1-Lipschitz, GCBS(D) implies by the classical Chernoff bound that for all $r \geq 0$ we have

$$\mu\{x \in \Omega : d(x_0, x) > \mu(d) + r\} \leq e^{-\frac{r^2}{4D}} \quad (36)$$

where

$$\mu(d) := \int d(x_0, x) \, d\mu(x).$$

We have

$$\begin{aligned} & \int e^{ad(x_0, x)^2} \, d\mu(x) \\ &= \int e^{ad(x_0, x)^2} \mathbb{1}_{\{d(x, x_0) < \mu(d)\}} \, d\mu(x) + \int e^{ad(x_0, x)^2} \mathbb{1}_{\{d(x, x_0) \geq \mu(d)\}} \, d\mu(x) \\ &\leq e^{a\mu(d)^2} + e^{2a\mu(d)^2} \int e^{2a(d(x_0, x) - \mu(d))^2} \mathbb{1}_{\{d(x, x_0) \geq \mu(d)\}} \, d\mu(x). \end{aligned}$$

Now we use the fact that

$$\begin{aligned} & \int e^{2a(d(x_0, x) - \mu(d))^2} \mathbb{1}_{\{d(x, x_0) \geq \mu(d)\}} \, d\mu(x) \\ &= 1 + \int_1^\infty \mu\left(\left\{x : e^{2a(d(x_0, x) - \mu(d))^2} \geq u\right\}\right) \, du. \end{aligned}$$

The result follows using (36) with $a = 1/(16D)$.

Statement 2. Since for all x and for all $a > 0$

$$d(x_0, x) \leq \frac{1}{\sqrt{a}} e^{ad(x_0, x)^2}$$

it follows that $x \mapsto d(x_0, x)$ is μ -integrable. We also have that e^f is μ -integrable for any Lipschitz function. Now, using Jensen's inequality and then the triangle inequality, we obtain

$$\begin{aligned} & \int e^{f - \mu(f)} \, d\mu \leq \int \int e^{f(x) - f(y)} \, d\mu(x) \, d\mu(y) \\ &\leq \int \int e^{\text{lip}(f)d(x, y)} \, d\mu(x) \, d\mu(y) \leq \left(\int e^{\text{lip}(f)d(x, x_0)} \, d\mu(x) \right)^2. \quad (37) \end{aligned}$$

Combining the elementary inequality

$$\text{lip}(f) d(x, x_0) \leq \frac{\text{lip}(f)^2}{4a} + ad(x, x_0)^2$$

with (24), we obtain

$$\int e^{\text{lip}(f)d(x, x_0)} d\mu(x) \leq b e^{\frac{1}{4a} \text{lip}(f)^2} .$$

This implies

$$\int \int e^{f(x)-f(y)} d\mu(x) d\mu(y) \leq b^2 e^{\frac{1}{2a} \text{lip}(f)^2} . \quad (38)$$

We now show that the pre-factor of the exponential can be changed to 1. We first establish the following lemma.

LEMMA A.1. *Let Z be a random variable with all odd moments vanishing and such that there exist $C_1 \geq 1$ and $C_2 > 0$ such that for all $\lambda \in \mathbb{R}$*

$$\mathbb{E} (e^{\lambda Z}) \leq C_1 e^{C_2 \lambda^2} .$$

Then for all $\lambda \in \mathbb{R}$ we have

$$\mathbb{E} (e^{\lambda Z}) \leq e^{C_2 \left(1 \vee \frac{C_1 e}{2\sqrt{\pi}}\right) \lambda^2} .$$

PROOF. We have for any $\lambda \in \mathbb{R}$

$$\mathbb{E}(Z^{2q}) = \mathbb{E}(Z^{2q} e^{-\lambda Z} e^{\lambda Z}) \leq C_1 (2q)^{2q} \lambda^{-2q} e^{-2q} e^{C_2 \lambda^2} \leq C_1 4^q q^q e^{-q} C_2^q$$

where the first inequality follows by maximizing $x^{2q} e^{-\lambda x}$ over x , while the second is obtained by minimizing over λ . Using the bound

$$\sqrt{2\pi} n^{n+\frac{1}{2}} e^{-n} \leq n! \leq e n^{n+\frac{1}{2}} e^{-n}$$

which is valid for all $n \geq 1$, we get

$$\frac{C_1 4^q q^q e^{-q} C_2^q}{(2q)!} \leq \frac{\left(1 \vee \frac{C_1 e}{2\sqrt{\pi}}\right)^q C_2^q}{q!}, \quad q \geq 1 .$$

The result follows. \square

We now apply the above lemma to the random variable $Z = f(X) - f(Y)$, where (X, Y) is distributed according to the product probability measure $d\mu(x) d\mu(y)$. It is easy to verify that all odd moments vanish (Z is anti-symmetrical with respect to the exchange of x and y) and the bound on the exponential moments follow by replacing f by λf in (38). We use the constants $C_1 = b^2$ and $C_2 = \frac{\text{lip}(f)^2}{2a}$. The second part of Theorem 5.1 follows from the first inequality in (37).

B An approximation lemma

In this appendix, $(\Omega, \|\cdot\|)$ is a separable Banach space. We denote by $\text{Lip}(\Omega, \mathbb{R})$ the space of real-valued Lipschitz functions on $(\Omega, \|\cdot\|)$, by $\text{Lip}_s(\Omega, \mathbb{R})$ the space of real-valued Lipschitz functions with bounded support, and by $\text{Lip}_b(\Omega, \mathbb{R})$ the space of real-valued bounded Lipschitz functions. We denote by $\mathcal{C}^\infty(\Omega, \mathbb{R})$ the space of real-valued infinitely differentiable functions, and by $\mathcal{C}_s^\infty(\Omega, \mathbb{R})$ the space of real-valued infinitely differentiable functions with bounded support.

Let \mathcal{C} be a class of real-valued functions on Ω . We say that μ satisfies $\text{GCB}(\mathcal{C}; D)$ if there exists $D > 0$ such that

$$\log \mu \left(e^{f - \mu(f)} \right) \leq D \text{lip}(f)^2$$

for all $f \in \mathcal{C}$.

LEMMA B.1. *Let μ be a probability measure on Ω . Then*

1. *If μ satisfies $\text{GCB}(\mathcal{C}_s^\infty(\Omega, \mathbb{R}); D)$ then it satisfies $\text{GCB}(\text{Lip}_s(\Omega, \mathbb{R}); D)$.*
2. *If μ satisfies $\text{GCB}(\text{Lip}_s(\Omega, \mathbb{R}); D)$ then it satisfies $\text{GCB}(\text{Lip}(\Omega, \mathbb{R}); D)$.*

PROOF. Let ν be a \mathcal{C}^∞ (in the sense of distributions) probability measure on Ω with bounded support. For every $\lambda > 0$ we define the rescaled measure ν_λ by

$$\nu_\lambda(f) := \nu(f_\lambda)$$

for any f continuous with bounded support, where $f_\lambda(x) := f(\lambda x)$. For $f \in \text{Lip}_s(\Omega, \mathbb{R})$, we have $\nu_\lambda * f \in \mathcal{C}_s^\infty(\Omega, \mathbb{R})$ and $\text{lip}(\nu_\lambda * f) \leq \text{lip}(f)$. Since μ is assumed to satisfy $\text{GCB}(\mathcal{C}_s^\infty(\Omega, \mathbb{R}); D)$, it follows that

$$\mu \left(e^{\nu_\lambda * f - \mu(\nu_\lambda * f)} \right) \leq e^{D \text{lip}(f)^2}.$$

The first statement then follows by dominated convergence.

For the second statement, as an intermediate step, we prove that if μ satisfies $\text{GCB}(\text{Lip}_s(\Omega, \mathbb{R}); D)$ then it satisfies $\text{GCB}(\text{Lip}_b(\Omega, \mathbb{R}); D)$. Let $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be defined by

$$\psi(u) = \begin{cases} 1 & \text{if } u \leq 1 \\ 2 - u & \text{if } 1 \leq u \leq 2 \\ 0 & \text{if } u \geq 2. \end{cases}$$

For any $A > 0$ define $\psi_A : \Omega \rightarrow \mathbb{R}^+$ by

$$\psi_A(x) = \psi \left(\frac{\|x\|}{A} \right).$$

We have $\psi_A \in \text{Lip}_s(\Omega, \mathbb{R})$ and $\text{lip}(\psi_A) \leq 1/A$. Take $f \in \text{Lip}_b(\Omega, \mathbb{R})$ such that $f(0) = 0$ (without loss of generality), define the function F_A by

$$F_A(x) = f(x)\psi_A(x).$$

We show that $F_A \in \text{Lip}_s(\Omega, \mathbb{R})$. We have

$$F_A(x) - F_A(y) = f(x)[\psi_A(x) - \psi_A(y)] + \psi_A(y)[f(x) - f(y)].$$

Since $\|\psi_A\|_\infty \leq 1$ we get

$$\text{lip}(F_A) \leq \frac{\|f\|_\infty}{A} + \text{lip}(f).$$

Since μ is assumed to satisfy $\text{GCB}(\text{Lip}_s(\Omega, \mathbb{R}); D)$, we have

$$\mu(e^{F_A - \mu(F_A)}) \leq \exp\left(D\left(\frac{\|f\|_\infty}{A} + \text{lip}(f)\right)^2\right).$$

Using the Dominated Convergence Theorem, we take the limit $A \rightarrow +\infty$ and get

$$\mu(e^{f - \mu(f)}) \leq e^{D \text{lip}(f)^2}.$$

Finally, let us prove that if μ satisfies $\text{GCB}(\text{Lip}_b(\Omega, \mathbb{R}); D)$ then it satisfies $\text{GCB}(\text{Lip}(\Omega, \mathbb{R}); D)$. Define for $M > 0$

$$f_M(x) = (f(x) \wedge M) \vee (-M).$$

By observing that $\text{lip}(f_M) \leq \text{lip}(f)$ and since μ satisfies $\text{GCB}(\text{Lip}_b(\Omega, \mathbb{R}); D)$ by assumption, we have

$$\mu(e^{f_M - \mu(f_M)}) \leq e^{D \text{lip}(f)^2}. \quad (39)$$

We are going to take the limit $M \rightarrow +\infty$ and prove that the left-hand side converges to $\mu(\exp(f - \mu(f)))$. We first prove that $\sup_{M>0} |\mu(f_M)| < +\infty$. We start by proving that $\inf_{M>0} \mu(f_M) > -\infty$. Take a ball B such that $\mu(B) > 0$. Denote by x_B its center and by r_B its radius. Using (39) and the mean-value theorem, we deduce that there exists $y_M \in B$ such that

$$\mu(B) e^{f_M(y_M) - \mu(f_M)} \leq e^{D \text{lip}(f)^2}.$$

Hence, using that $\text{lip}(f_M) \leq \text{lip}(f)$, we get

$$f_M(x_B) \leq \mu(f_M) + D \text{lip}(f)^2 - \log \mu(B) + \text{lip}(f)r_B.$$

Since $f_M(0) = 0$, we obtain $f_M(x_B) \geq -\text{lip}(f)\|x_B\|$, which implies $\inf_{M>0} \mu(f_M) > -\infty$. A similar argument applies to $-f$, therefore

$$A_f := \sup_{M>0} |\mu(f_M)| < +\infty.$$

We now prove that e^f is integrable with respect to μ . We have

$$\mu(e^{f_M}) = \mu(\mathbf{1}_{\{f \geq 0\}} e^{f_M}) + \mu(\mathbf{1}_{\{f < 0\}} e^{f_M}). \quad (40)$$

If $x \in \Omega$ is such that $f(x) \geq 0$, then $f_M(x) \uparrow f(x)$ as $M \uparrow +\infty$, then

$$\mu(\mathbf{1}_{\{f \geq 0\}} e^{f_M}) \leq \mu(e^{f_M}) \leq e^{D \text{lip}(f)^2 + A_f}.$$

By the Monotone Convergence Theorem we thus get

$$\mu(\mathbf{1}_{\{f \geq 0\}} e^f) = \lim_{M \rightarrow +\infty} \mu(\mathbf{1}_{\{f \geq 0\}} e^{f_M}) \leq e^{D \text{lip}(f)^2 + A_f}. \quad (41)$$

Now we deal with the second term in the right-hand side of (40). Since the function $\mathbf{1}_{\{f < 0\}} e^{f_M}$ is nonnegative and bounded above by 1 and converges pointwise to $\mathbf{1}_{\{f < 0\}} e^f$ as M tends to $+\infty$, we apply the Dominated Convergence Theorem to get that

$$\lim_{M \rightarrow +\infty} \mu(\mathbf{1}_{\{f < 0\}} e^{f_M}) = \mu(\mathbf{1}_{\{f < 0\}} e^f).$$

Therefore, using this inequality, (41) and (40) we conclude that

$$\lim_{M \rightarrow +\infty} \mu(e^{f_M}) = \mu(e^f) < +\infty. \quad (42)$$

By a similar argument one shows that $\mu(e^{-f}) < +\infty$.

We now prove that $\mu(f_M)$ converges to $\mu(f)$ as M tends to $+\infty$. We observe that $|f_M| \leq e^f + e^{-f}$. Hence by the Dominated Convergence Theorem we conclude that

$$\lim_{M \rightarrow +\infty} \mu(f_M) = \mu(f). \quad (43)$$

Using (43) and (42), we can take the limit $M \rightarrow +\infty$ in inequality (39) and obtain

$$\mu(e^{f-\mu(f)}) \leq e^{D \text{lip}(f)^2}.$$

The lemma is proved. \square

Acknowledgment

F. Redig thanks Ecole Polytechnique and the CNRS for financial support and hospitality. We thank R. Kraaij for useful discussions in an early stage of this work.

References

- [1] D. Bakry, I. Gentil, M. Ledoux. *Analysis and geometry of Markov diffusion operators*. Grundlehren der Mathematischen Wissenschaften, 348. Springer, 2014.
- [2] V. Bansaye, P. Collet, S. Martínez, S. Méléard, J. San Martín. Diffusions from infinity. *Trans. AMS* **372**, 57815823 (2019).
- [3] S. Boucheron, G. Lugosi, P. Massart. *Concentration inequalities. A nonasymptotic theory of independence*. Oxford University Press, 2013.
- [4] J.-R. Chazottes, P. Collet, F. Redig. On concentration inequalities and their applications for Gibbs measures in lattice systems. *J. Stat. Phys.* 169 (2017), no. 3, 504–546.
- [5] J.-R. Chazottes, P. Collet, C. Külske, F. Redig. Concentration inequalities for random fields via coupling. *Probab. Theory Related Fields* 137 (2007), no. 1-2, 201225.
- [6] B. Davis. On the L_p norms of stochastic integrals and other martingales, *Duke Math. J.* 43, 697-704 (1976).
- [7] A.C.D. van Enter, R. Fernández, F. den Hollander, F. Redig. Possible loss and recovery of Gibbsianness during the stochastic evolution of Gibbs measures. *Comm. Math. Phys.* 226 (2002), no. 1, 101130.
- [8] H. Djellout, A. Guillin, and L. Wu. Transportation cost-information inequalities and applications to random dynamical systems and diffusions. *Ann. Probab.*, Volume 32, Number 3B (2004), 2702-2732.
- [9] M.Z. Guo, G.C. Papanicolaou, S.R.S. Varadhan, Nonlinear diffusion limit for a system with nearest neighbor interactions. *Comm. Math. Phys.* 118 (1988), no. 1, 31–59.
- [10] M. Hirsch, S. Smale, R. Devaney. *Differential equations, dynamical systems, and an introduction to chaos*. Second edition. Pure and Applied

Mathematics (Amsterdam), 60. Elsevier/Academic Press, Amsterdam, 2004.

- [11] C. Külske. Concentration inequalities for functions of Gibbs fields with application to diffraction and random Gibbs measures. *Comm. Math. Phys.* 239 (2003), no. 1-2, 29-51.
- [12] M. Ledoux. *The concentration of measure phenomenon*. Mathematical Surveys and Monographs, 89. American Mathematical Society, 2001.
- [13] P. Mattila. *Geometry of Sets and Measures in Euclidean Spaces*. Cambridge University Press, 1995.
- [14] M.K. von Renesse, K.T. Sturm, Transport inequalities, gradient estimates, entropy and Ricci curvature, *Comm. Pure and Applied Math.*, LVIII, (2005), 0923-0940,