Multiparameter scaling of the Kondo effect in quantum dots with an even number of electrons

Mikio Eto¹ and Yuli V. Nazarov²

¹*Faculty of Science and Technology, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan*

²*Department of Applied Physics/DIMES, Delft University of Technology, Lorentzweg 1, 2628 CJ Delft, The Netherlands*

(Received 7 March 2002; published 31 October 2002)

We address a recent theoretical discrepancy concerning the Kondo effect in quantum dots with an even number of electrons where spin-singlet and -triplet states are nearly degenerate. We show that the discrepancy arises from the fact that the Kondo scaling involves many parameters, which makes the results depend on concrete microscopic models. We illustrate this by the scaling calculations of the Kondo temperature T_K as a function of the energy difference between the singlet and triplet states Δ . $T_K(\Delta)$ decreases with increasing Δ , showing a crossover from a power law with a universal exponent to that with a nonuniversal exponent. The crossover depends on the initial parameters of the model.

DOI: 10.1103/PhysRevB.66.153319

PACS number(s): 73.23.Hk, 72.15.Qm, 85.35.Gv

Recent observations of the Kondo effect in semiconductor quantum dots have attracted much attention.¹ In quantum dots, the number of electrons, N , is fixed by the Coulomb blockade to integer values and can be tuned by gate voltages. The usual Kondo effect takes place for an odd N . Here, the spin 1/2 is formed in the dot and it is coupled to the Fermi sea of conduction electrons in external leads through tunneling barriers.

Strikingly the Kondo effect has also been observed for an even N in some special cases.²⁻⁴ In so-called “vertical” quantum dots, the ground state can be changed from a spin triplet to singlet by the magnetic field.⁵ Sasaki *et al.* have found a pronounced Kondo effect near the degeneracy point between the spin states.² In this experiment the Zeeman splitting of the triplet state can be neglected owing to a small g factor in semiconductor heterostructures.⁶

In our previous paper, we have suggested an explanation of this phenomenon within a model.⁷ The Kondo temperature T_K has been evaluated as a function of the energy difference between the singlet and triplet states, Δ , using the “poor man’s” scaling method. We have shown that (i) $T_K(\Delta)$ is maximal around $\Delta=0$, (ii) for positive Δ , $T_K(\Delta)$ decreases with increasing Δ obeying a power law, $T_K(\Delta) \propto 1/\Delta^\gamma$, and (iii) for negative Δ , the Kondo effect is not relevant when $|\Delta| \gg T_K(0)$. The exponent γ is not universal ($0 < \gamma \leq 1$) depending on a ratio of the initial coupling constants. Our results have indicated an enhancement of the Kondo effect by the competition between singlet and triplet states, qualitatively in agreement with the experimental results by Sasaki *et al.*² After our work, Pustilnik and Glazman have considered a different model for this “triplet-singlet Kondo effect.”⁸ Their results are qualitatively the same as ours but quantitatively different: They have derived a power law of $T_K(\Delta)$ with a universal exponent, $\gamma=2+\sqrt{5}$.

The purpose of this brief report is to elucidate the discrepancy between the results mentioned above. It should be noted that a microscopic model sets the initial values of coupling constants, which are different between Refs. 7 and 8, whereas a common system of the scaling equations describes the evolution of the coupling constants. If it were a one-parameter scaling, both models would result in the identical

results. However, this is not the present case of *multiparameter* scaling. The scaling equations are

$$\begin{cases} dJ^{(1)}/d \ln D &= -2\nu[J^{(1)2} + (\tilde{J}_1^2 + \tilde{J}_2^2)/2], \\ dJ^{(2)}/d \ln D &= -2\nu[J^{(2)2} + (\tilde{J}_1^2 + \tilde{J}_2^2)/2], \\ d\tilde{J}_1/d \ln D &= -2\nu(J^{(1)} + J^{(2)})\tilde{J}_1 + \nu J' \tilde{J}_1, \\ d\tilde{J}_2/d \ln D &= -2\nu(J^{(1)} + J^{(2)})\tilde{J}_2 - \nu J' \tilde{J}_2, \\ dJ'/d \ln D &= 8\nu(\tilde{J}_1^2 - \tilde{J}_2^2), \end{cases} \quad (1)$$

when the energy scale D is much larger than $|\Delta|$. This system of the equations encompasses five coupling constants ($J^{(1)}, J^{(2)}, \tilde{J}_1, \tilde{J}_2, J'$) and exhibits a complicated nonlinear behavior. Our previous results correspond to an unstable fixed line of these equations, whereas the results of Ref. 8 to a stable fixed line. We show that, in general, the coupling constants develop along the unstable fixed line at first and transit toward the stable fixed line with decreasing D . The stable fixed line is much less relevant than in the situation of one-parameter scaling. Indeed our numerical calculations indicate that it may take up to six decades to reach the neighborhood of the stable fixed line [see Fig. 1(a)]. We find that $T_K(\Delta)$ exhibits a crossover between power laws with nonuniversal and universal exponents, which are determined by the unstable and stable fixed lines, respectively. Besides, we show that our previous calculations along the unstable fixed line provide a lower limit of $T_K(\Delta)$ in an analytical form.

In a Coulomb blockade region with an even N , we consider two electrons at the background of a singlet state of the other $N-2$ electrons. We regard the background as the vacuum $|0\rangle$. The two electrons occupy two levels of different orbital symmetry, $\varepsilon_1, \varepsilon_2$, and make a spin-triplet ($S=1$) or -singlet state ($S=0$): $|SM\rangle = |11\rangle = d_{1\uparrow}^\dagger d_{2\uparrow}^\dagger |0\rangle$, $|10\rangle = (1/\sqrt{2})(d_{1\uparrow}^\dagger d_{2\downarrow}^\dagger + d_{1\downarrow}^\dagger d_{2\uparrow}^\dagger) |0\rangle$, $|1-1\rangle = d_{1\downarrow}^\dagger d_{2\downarrow}^\dagger |0\rangle$, $|00\rangle = (1/\sqrt{2})(C_1 d_{1\uparrow}^\dagger d_{1\downarrow}^\dagger - C_2 d_{2\uparrow}^\dagger d_{2\downarrow}^\dagger) |0\rangle$, where $d_{i\sigma}^\dagger$ creates an electron with spin σ in level i . The energy difference between the spin states is denoted by $\Delta = E_{S=0} - E_{S=1}$. The coefficients in the singlet state, C_1 and C_2 ($|C_1|^2 + |C_2|^2 = 2$), should be determined by the electron-electron interaction and one-electron level spacing, $\delta = \varepsilon_2 - \varepsilon_1$.⁹ We have set

FIG. 1. The scaling of the coupling constants by Eqs. (1), assuming $\Delta = 0$. (a) $(J^{(1)} - J^{(2)})/(J^{(1)} + J^{(2)})$ and (b) $(\tilde{J}_1 - \tilde{J}_2)/(J^{(1)} + J^{(2)})$ as functions of $\ln D/T_K(0)$, where $T_K(0)$ is the Kondo temperature at $\Delta = 0$. $J^{(2)}/J^{(1)}$ is *a*, 1; *b*, 0.5; and *c*, 0.3. The cases with $C_1 = C_2$ are represented by solid lines, $C_2 = 0$ by broken lines, and $C_2/C_1 = 0.6$ by dotted lines. Note that three lines overlap for case *a* in (a). Initially, we choose $\nu(J^{(1)} + J^{(2)}) = 0.01$.

$C_1 = C_2$ in our previous work,⁷ whereas $C_2 = 0$ in Ref. 8. Now we investigate general cases with respect to C_1 and C_2 . The other singlet states with higher energies are disregarded.

The level i ($= 1, 2$) in the dot is connected to two external leads L, R , by the tunnel couplings $V_{L,i}, V_{R,i}$. We assume two channels in the leads; the orbital symmetry is conserved in the tunneling processes.¹⁰ We perform the unitary transformation for the conduction electrons in the two leads,¹¹ $c_{k\sigma}^{(i)} = (V_{L,i}^* c_{L,k\sigma}^{(i)} + V_{R,i}^* c_{R,k\sigma}^{(i)})/V_i$, $\bar{c}_{k\sigma}^{(i)} = (-V_{R,i} c_{L,k\sigma}^{(i)} + V_{L,i} c_{R,k\sigma}^{(i)})/V_i$ with $V_i = \sqrt{|V_{L,i}|^2 + |V_{R,i}|^2}$, where $c_{\alpha,k\sigma}^{(i)}$ is the annihilation operator of an electron in lead α , with momentum k , spin σ , and orbital symmetry i ($= 1, 2$). The mode $c_{k\sigma}^{(i)}$ is coupled to the dot with V_i . The mode $\bar{c}_{k\sigma}^{(i)}$ is decoupled from the leads, which shall be disregarded. The Hamiltonian of the leads and that of tunneling processes are written as

$$H_{\text{leads}} = \sum_{k\sigma i} \epsilon_k^{(i)} c_{k\sigma}^{(i)\dagger} c_{k\sigma}^{(i)}, \quad (2)$$

$$H_T = \sum_{k\sigma i} V_i (c_{k\sigma}^{(i)\dagger} d_{i\sigma} + \text{H.c.}), \quad (3)$$

respectively. The density of states ν in the leads remains constant in the energy band of $[-D, D]$.

In the Coulomb blockade region, the addition and extraction energies, $E^\pm \equiv E(N \pm 1) - E(N) \mp \mu$, are much larger than the level broadening $\Gamma_i = \pi \nu V_i^2$ ($i = 1, 2$) and temperature T , where $E(N)$ is the energy with N electrons in the dot and μ is the Fermi energy in the leads. We also assume that

$E^\pm \gg |\Delta|, \delta$. As in Ref. 7, we integrate out the states with $N \pm 1$ electrons and obtain the effective low-energy Hamiltonian,

$$H_{\text{eff}} = H_{\text{leads}} + H_{\text{dot}} + H^{S=1} + H^{S=1 \leftrightarrow 0} + H'_{\text{eff}}. \quad (4)$$

The Hamiltonian of the dot, H_{dot} , reads

$$H_{\text{dot}} = \sum_{S,M} E_S f_{SM}^\dagger f_{SM}, \quad (5)$$

using pseudofermion operators $f_{SM}^\dagger$ (f_{SM}) that create (annihilate) the state $|SM\rangle$. It is required that $\sum_{S,M} f_{SM}^\dagger f_{SM} = 1$. $H^{S=1}$ describes the spin-flip processes among three components of the triplet state,

$$H^{S=1} = \sum_{kk'} \sum_{i=1,2} J^{(i)} [\sqrt{2} (f_{11}^\dagger f_{10} + f_{10}^\dagger f_{1-1}) c_{k'\uparrow}^{(i)\dagger} c_{k\uparrow}^{(i)} + \sqrt{2} (f_{10}^\dagger f_{11} + f_{1-1}^\dagger f_{10}) c_{k'\uparrow}^{(i)\dagger} c_{k\downarrow}^{(i)} + (f_{11}^\dagger f_{11} - f_{1-1}^\dagger f_{1-1}) \times (c_{k'\uparrow}^{(i)\dagger} c_{k\uparrow}^{(i)} - c_{k'\downarrow}^{(i)\dagger} c_{k\downarrow}^{(i)})]. \quad (6)$$

The exchange couplings are $J^{(i)} = V_i^2/(2E_c)$ ($i = 1, 2$), with $1/E_c = 1/E^+ + 1/E^-$, which are accompanied by the scattering of a conduction electron of channel i . $H^{S=1 \leftrightarrow 0}$ represents the conversion between the triplet and singlet states, with the interchannel scattering of a conduction electron,

$$H^{S=1 \leftrightarrow 0} = \sum_{kk'} [\sqrt{2} (\tilde{J}_1 f_{11}^\dagger f_{00} - \tilde{J}_2 f_{00}^\dagger f_{1-1}) c_{k'\uparrow}^{(1)\dagger} c_{k\uparrow}^{(2)} + \sqrt{2} (\tilde{J}_2 f_{00}^\dagger f_{11} - \tilde{J}_1 f_{1-1}^\dagger f_{00}) c_{k'\uparrow}^{(1)\dagger} c_{k\downarrow}^{(2)} - (\tilde{J}_1 f_{10}^\dagger f_{00} + \tilde{J}_2 f_{00}^\dagger f_{10}) (c_{k'\uparrow}^{(1)\dagger} c_{k\uparrow}^{(2)} - c_{k'\downarrow}^{(1)\dagger} c_{k\downarrow}^{(2)}) + (1 \leftrightarrow 2)], \quad (7)$$

where $\tilde{J}_i = C_i V_i V_2/(2E_c)$. The last term of H_{eff} represents the scattering processes without spin flip in the dot,

$$H'_{\text{eff}} = \sum_{kk'\sigma} \sum_{i=1,2} \left[J'^{(i)} c_{k'\sigma}^{(i)\dagger} c_{k\sigma}^{(i)} \sum_M f_{1M}^\dagger f_{1M} + J''^{(i)} c_{k'\sigma}^{(i)\dagger} c_{k\sigma}^{(i)} f_{00}^\dagger f_{00} \right]. \quad (8)$$

We calculate T_K using the poor man's scaling method.¹² By changing the energy scale (bandwidth of the conduction electrons) from D to $D - |dD|$, we renormalize the exchange couplings not to change the low-energy physics, within the second-order perturbation with respect to $H^{S=1} + H^{S=1 \leftrightarrow 0} + H'_{\text{eff}}$. This procedure yields the scaling equations in two limits. For $D \gg |\Delta|$, H_{dot} can be safely disregarded in H_{eff} . The scaling equations are given by Eqs. (1) with $J' = J'^{(1)} - J'^{(2)} - J''^{(1)} + J''^{(2)}$. For $D \ll \Delta$, the ground state of the dot is a spin triplet and the singlet state can be disregarded. Then $J^{(1)}$ and $J^{(2)}$ evolve independently,

$$\frac{d}{d \ln D} J^{(i)} = -2\nu J^{(i)2}, \quad (9)$$

whereas the other coupling constants do not change. The Kondo temperature is determined as the energy scale at which the coupling constants become so large that the perturbation breaks down.

In our previous model,⁷ $\tilde{J}_1 = \tilde{J}_2$ and the scaling equations (1) are closed without J' . The ratio of $J^{(2)}/J^{(1)}$ is also fixed in the scaling by Eqs. (1). We obtain the Kondo temperature as a function of Δ , as follows.

(i) When $|\Delta| \ll T_K(0)$, the scaling equations (1) remain valid till the scaling ends. This yields

$$T_K(0) = D_0 \exp[-1/2\nu(J^{(1)} + J^{(2)})], \quad (10)$$

where D_0 is the initial bandwidth, which is given by $\sqrt{E^+ E^-}$.¹³

(ii) When $\Delta > D_0$, the scaling equations (9) work in the whole scaling region. T_K is identical to that of a localized spin with $S=1$,¹⁴

$$T_K(\infty) = D_0 \exp[-1/2\nu J^{(1)}], \quad (11)$$

when $J^{(1)} \geq J^{(2)}$.

(iii) In the intermediate region of $T_K(0) \ll \Delta \ll D_0$, we match the solutions of Eqs. (1) and (9) at $D \approx \Delta$ and obtain a power law

$$T_K(\Delta) = T_K(0) [T_K(0)/\Delta]^\gamma. \quad (12)$$

The exponent is given by $\gamma = J^{(2)}/J^{(1)}$.

(iv) For $\Delta < 0$, T_K drops to zero suddenly at $\Delta \sim -T_K(0)$.

In general situations of $C_1 \neq C_2$, we find that the line of $J^{(2)}/J^{(1)} = \text{const}$ and $\tilde{J}_1 = \tilde{J}_2$ is not stable in the scaling by Eqs. (1). The renormalization flow finally goes to a fixed point of $J^{(1)} = J^{(2)} = \infty$, $J^{(2)}/J^{(1)} = 1$, $\tilde{J}_1/J^{(1)} = \sqrt{2(2+\sqrt{5})}$, $\tilde{J}_2/J^{(1)} = 0$, and $J'/J^{(1)} = -2(1+\sqrt{5})$, as the energy scale D decreases to T_K .¹⁵ Around this fixed point, the stable fixed line discussed in Ref. 8 is relevant. By expanding the coupling constants around the fixed point to the first order of $1/\ln(D/T_K)$, Pustilnik and Glazman have obtained a power law of $T_K(\Delta)$ with a universal exponent, $\gamma = 2 + \sqrt{5}$.⁸ However, the range of energy scale is very limited where the physical properties are determined by the stable fixed line, due to the complexity of the multiparameter scaling.

To elucidate this multiparameter scaling, we solve the scaling equations numerically when $\Delta \geq 0$; Eqs. (1) for $D > \Delta$ and Eqs. (9) for $D < \Delta$. The Kondo temperature T_K is determined as the energy scale D at which $\nu(J^{(1)} + J^{(2)}) = 1$. First we examine the scaling by Eqs. (1), assuming $\Delta = 0$. In Fig. 1(a), we show the ratio of $(J^{(1)} - J^{(2)})/(J^{(1)} + J^{(2)})$ as a function of $\ln D/T_K(0)$, where $T_K(0)$ is the Kondo temperature at $\Delta = 0$. We set $J^{(2)}/J^{(1)}$ to be a , 1; b , 0.5; and c , 0.3 in the initial condition.¹⁶ In the case of $C_1 = C_2$ (solid lines), the ratio is fixed at a nonuniversal value determined by the given condition. In the cases of $C_2 = 0$ (broken lines) and $C_2/C_1 = 0.6$ (dotted lines), the ratio gradually deviates from the initial value with decreasing D ,¹⁷ and finally goes to zero. This limit corresponds to the fixed point of the renormalization, $J^{(2)}/J^{(1)} = 1$. It may need to reduce the energy scale by six decades to reach its vicinity.

FIG. 2. The Kondo temperature, $T_K/T_K(0)$, as a function of $\Delta/T_K(0)$, on a log-log scale ($\Delta > 0$). $T_K(0)$ is the Kondo temperature at $\Delta = 0$. $J^{(2)}/J^{(1)}$ is a , 1, and b , 0.3. The cases with $C_1 = C_2$ are represented by solid lines, $C_2 = 0$ by broken lines, and $C_2/C_1 = 0.6$ by dotted lines. Initially, we choose $\nu(J^{(1)} + J^{(2)}) = 0.01$.

(The case a with $J^{(1)} = J^{(2)}$ is an exception, where the fixed point is on the line of $J^{(2)}/J^{(1)} = \text{const}$.) Similarly Fig. 1(b) indicates $(\tilde{J}_1 - \tilde{J}_2)/(J^{(1)} + J^{(2)})$. In the case of $C_1 = C_2$, the ratio is fixed at zero ($\tilde{J}_1 = \tilde{J}_2$). Otherwise, the ratio goes to a value of the fixed point, $\sqrt{2 + \sqrt{5}}/\sqrt{2}$, as D decreases.

Figure 2 presents the Kondo temperature as a function of Δ on a log-log scale; a , $J^{(2)}/J^{(1)} = 1$ and b , 0.3. Both T_K and Δ are normalized by $T_K(0)$. In all the cases, the Kondo temperature increases with decreasing Δ . In the case of $C_1 = C_2$ (solid lines), $T_K(\Delta)/T_K(0)$ obeys a power law, Eq. (12), with a nonuniversal exponent, $\gamma = J^{(2)}/J^{(1)}$. In the cases of $C_2 = 0$ (broken lines) and $C_2/C_1 = 0.6$ (dotted lines), $T_K(\Delta)$ shows a crossover from the power law with $\gamma = J^{(2)}/J^{(1)}$ at large $\Delta/T_K(0)$ to that with $\gamma = 2 + \sqrt{5} \approx 4.2$ at

FIG. 3. The Kondo temperature T_K as a function of Δ ($\Delta > 0$). The unit of T_K is $D_0 \exp[-1/2\nu(J^{(1)} + J^{(2)})]$, where D_0 is the initial bandwidth ($D_0 = \sqrt{E^+ E^-} \approx 5$ K in the experiment of Ref. 2). $J^{(2)}/J^{(1)}$ is a , 1; b , 0.5; and c , 0.3. The cases with $C_1 = C_2$ are represented by solid lines, $C_2 = 0$ by broken lines, and $C_2/C_1 = 0.6$ by dotted lines. As the initial condition, we choose $\nu(J^{(1)} + J^{(2)}) = 0.1$. Inset: The Kondo temperature at $\Delta = 0$, $T_K(0)$, as a function of θ , where $\tan^2 \theta = J^{(2)}/J^{(1)}$. $\theta/\pi = 0.25$ (0.20, 0.15) for $J^{(2)}/J^{(1)} = 1$ (0.5, 0.3).

small $\Delta/T_K(0)$. The crossover depends on the initial values of C_2/C_1 and $J^{(2)}/J^{(1)}$. The universal exponent is seen in quite limited situations.

Finally we compare the Kondo temperature with various C_2/C_1 and $J^{(2)}/J^{(1)}$ in the initial condition. We choose $\nu(J^{(1)}+J^{(2)})=0.1$ which corresponds to the experimental situation with $E^\pm \approx 5$ K and level broadening $\Gamma \approx 1.5$ K.² In Fig. 3, we show T_K in units of $D_0 \exp[-1/2\nu(J^{(1)}+J^{(2)})]$ [which is identical to $T_K(0)$ with $C_1=C_2$, Eq. (10)], as a function of Δ . We find that the case with $C_1=C_2$ (solid lines) provides a lower limit of T_K for each value of $J^{(2)}/J^{(1)}$ ($a, 1$; $b, 0.5$; $c, 0.3$), whereas the case with $C_2=0$ (broken lines) gives an upper limit. The inset to Fig. 3 shows $T_K(0)$ as a function of $J^{(2)}/J^{(1)}$ ($\equiv \tan^2 \theta$). For fixed C_2/C_1 , $T_K(0)$ is maximal at $J^{(2)}/J^{(1)}=1$ ($\theta/\pi=0.25$) and decreases with decreasing $J^{(2)}/J^{(1)}$. In a case of $C_2=0$ and $J^{(2)}/J^{(1)}=1$, the development of the coupling constants by Eqs. (1) is restricted in a subspace with $\tilde{J}_2(D)=0$ and $J^{(1)}(D)=J^{(2)}(D)$, and in consequence they reach the fixed point fastest. This results in the highest T_K for a given value of $J^{(1)}+J^{(2)}$. As the initial values are deviated from the condition, the renormalization flow to the fixed point needs more

“time” in a larger space. [If D reaches Δ before T_K , the coupling constants follow Eqs. (9) at $T_K < D < \Delta$, which is common to all the cases.] Since our previous model with $C_1=C_2$ is the farthest from the condition of $C_2=0$, it yields a lower limit of the Kondo temperature.

In conclusion, we have examined a generalized model for the Kondo effect that involves spin-singlet and -triplet states in quantum dots to elucidate the discrepancy between the previous studies.^{7,8} The Kondo temperature T_K is calculated as a function of the energy difference between the states, Δ , using the scaling method. The function of $T_K(\Delta)$ shows a crossover between power laws with a nonuniversal exponent ($\gamma=J^{(2)}/J^{(1)}$) and with a universal exponent ($\gamma=2+\sqrt{5}$). Although our previous model with $\tilde{J}_1=\tilde{J}_2$ ($C_1=C_2$ in the singlet state)⁷ is not complete to discuss the triplet-singlet Kondo effect, it yields a lower limit of $T_K(\Delta)$ in an analytical form, Eqs. (10)–(12).

The authors gratefully acknowledge discussions with L. P. Kouwenhoven, S. De Franceschi, J. M. Elzerman, K. Majjala, S. Sasaki, W. G. van der Wiel, Y. Tokura, L. I. Glazman, M. Pustilnik, T. Saso, and G. E. W. Bauer.

¹D. Goldhaber-Gordon *et al.*, Nature (London) **391**, 156 (1998); D. Goldhaber-Gordon *et al.*, Phys. Rev. Lett. **81**, 5225 (1998); S.M. Cronenwett *et al.*, Science **281**, 540 (1998); F. Simmel *et al.*, Phys. Rev. Lett. **83**, 804 (1999); W.G. van der Wiel *et al.*, Science **289**, 2105 (2000).

²S. Sasaki *et al.*, Nature (London) **405**, 764 (2000).

³J. Schmid *et al.*, Phys. Rev. Lett. **84**, 5824 (2000).

⁴J. Nygård *et al.*, Nature (London) **408**, 342 (2000).

⁵S. Tarucha *et al.*, Phys. Rev. Lett. **77**, 3613 (1996).

⁶In carbon nanotubes, the Zeeman effect is much larger, which results in a different type of the Kondo effect with an even N (Ref. 4). This Kondo effect has been explained by M. Pustilnik *et al.*, Phys. Rev. Lett. **84**, 1756 (2000).

⁷M. Eto and Yu.V. Nazarov, Phys. Rev. Lett. **85**, 1306 (2000); M. Eto and Yu.V. Nazarov, Phys. Rev. B **64**, 085322 (2001).

⁸M. Pustilnik and L.I. Glazman, Phys. Rev. Lett. **85**, 2993 (2000); M. Pustilnik and L.I. Glazman, Phys. Rev. B **64**, 045328 (2001).

⁹ $C_1, C_2 \neq 0$ in general, owing to the matrix element of the Coulomb interaction between $d_{1\uparrow}^\dagger d_{1\downarrow}^\dagger |0\rangle$ and $d_{2\uparrow}^\dagger d_{2\downarrow}^\dagger |0\rangle$, $\langle 22|e^2/r|11\rangle \equiv K$. Note that $K=0$ in circular dots, if orbitals 1

and 2 have different angular momentums, which results in $C_2=0$.

¹⁰This is the case of vertical quantum dots in which the tunneling processes conserve the symmetry for the motion of electrons in the transverse direction.

¹¹L.I. Glazman and M.É. Raïkh, Pis'ma Zh. Éksp. Teor. Fiz. **47**, 378 (1988) [JETP Lett. **47**, 452 (1988)].

¹²P.W. Anderson, J. Phys. C **3**, 2436 (1970); P. Nozières and A. Blandin, J. Phys. (Paris) **41**, 193 (1980); D.L. Cox and A. Zawadowski, Adv. Phys. **47**, 599 (1998).

¹³F.D.M. Haldane, J. Phys. C **11**, 5015 (1978).

¹⁴I. Okada and K. Yosida, Prog. Theor. Phys. **49**, 1483 (1973).

¹⁵This has been discussed in Appendix C of the second paper in Ref. 8 in a special case of $J^{(1)}=J^{(2)}$.

¹⁶Initially, $\tilde{J}_i/(J^{(1)}+J^{(2)})=C_i\lambda/(1+\lambda^2)$ with $\lambda=\sqrt{J^{(2)}/J^{(1)}}$, and $J'/(J^{(1)}+J^{(2)})=(-C_1^2+C_2^2)/2$, assuming that $E^+=E^-$.

¹⁷At the beginning of the renormalization, $J^{(1)}$ and $J^{(2)}$ develop in the same way as in the case of $\tilde{J}_1=\tilde{J}_2$, to the lowest order of $\ln(D_0/D)$. To this order, $J^{(1)}/J^{(2)}$ is fixed and $T_K(\Delta)$ coincides with Eq. (12).