publication 2013.

Mare

CHICAGO JOURNALS

SHEA
The Society for Healthcare
Epidemiology of America

Improving Hand Hygiene Compliance in Hospitals by Design

Author(s): Marijke Melles, PhD; Vicki Erasmus, PhD; Martijn P. M. van Loon, MSc; Marc Tassoul, MSc; Ed F. van Beeck, MD, PhD; Margreet C. Vos, MD, PhD

Reviewed work(s):

Source: *Infection Control and Hospital Epidemiology*, Vol. 34, No. 1 (January 2013), pp. 102-103

Published by: [The University of Chicago Press](#) on behalf of [The Society for Healthcare Epidemiology of America](#)

Stable URL: <http://www.jstor.org/stable/10.1086/668772>

Accessed: 11/01/2013 01:55

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

The University of Chicago Press and The Society for Healthcare Epidemiology of America are collaborating with JSTOR to digitize, preserve and extend access to *Infection Control and Hospital Epidemiology*.

<http://www.jstor.org>

personal and programmatic insights to explain their proficiency and serve as models for their coworkers.

The primary limitation of our study is that we did not evaluate the impact of cleaning on surface contamination or healthcare-associated infections. Also, the study consisted of a small sample size of rooms and housekeepers, and the amount of time spent cleaning each room was not validated by direct observation. Although our findings relate to only a single setting, the thoroughness of cleaning that we documented is similar to that reported in many healthcare settings. If the efficiency of cleaning is similar in other settings, it is likely that substantial opportunities to improve both the thoroughness of cleaning and the overall efficiency of practice exist in many healthcare settings.

In conclusion, we documented a counter-intuitive observation that a greater amount of time spent cleaning a hospital room does not necessarily correlate with the effectiveness of cleaning high-touch surfaces. Our finding emphasizes that process improvement interventions should evaluate both the efficiency and thoroughness of hospital surface cleaning to optimize the cost effectiveness of cleaning practice in healthcare settings.

ACKNOWLEDGMENTS

We thank personnel in the Environmental Service Department for their dedicated service.

Potential conflicts of interest. M.E.R. reports receiving research support from 3M. P.C. reports serving as a consultant and on the speaker's bureau for Ecolab and Steris and having licensed patents to Ecolab. All other authors report no conflicts of interest relevant to this article. All authors submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest, and the conflicts that the editors consider relevant to this article are disclosed here.

**Mark E. Rupp, MD;¹ Ann Adler, RN;²
Margaret Schellen, BA;¹ Kyle Cassling, BA;¹
Teresa Fitzgerald, RN;² Lee Sholtz, RN;²
Elizabeth Lyden, MS;¹ Philip Carling, MD³**

Affiliations: 1. University of Nebraska Medical Center, Omaha, Nebraska; 2. Nebraska Medical Center, Omaha, Nebraska; 3. Boston University School of Medicine, Boston, Massachusetts.

Address correspondence to Mark Rupp, MD, 984031 Nebraska Medical Center, Omaha, NE 68198 (merupp@unmc.edu).

Received May 30, 2012; accepted September 5, 2012; electronically published November 20, 2012.

Presented in part: 5th Decennial International Conference on Healthcare Associated Infections 2010; Atlanta, Georgia; March 18–22, 2010.

Infect Control Hosp Epidemiol 2013;34(1):100–102

© 2012 by The Society for Healthcare Epidemiology of America. All rights reserved. 0899-823X/2013/3401-0017\$15.00. DOI: 10.1086/668779

REFERENCES

- Hayden MK, Bonten MJM, Blom DW, Lyle EA, van de Vijver DAMC, Weinstein RA. Reduction in acquisition of vancomycin-resistant enterococcus after enforcement of routine environmental cleaning measures. *Clin Infect Dis* 2006;42:1552–1560.
- Datta R, Platt R, Yokoe DS, Huang SS. Environmental cleaning intervention and risk of acquiring multidrug-resistant organisms from prior room occupants. *Arch Intern Med* 2011;171:491–494.
- Drees M, Snyderman DR, Schmid CH, et al. Prior environmental contamination increases the risk of acquisition of vancomycin-resistant enterococci. *Clin Infect Dis* 2008;46: 678–685.
- Huang SS, Datta R, Platt R. Risk of acquiring antibiotic-resistant bacteria from prior room occupants. *Arch Intern Med* 2006;166: 1945–1951.
- Dancer SJ, White LE, Lamb J, Girvan EK, Robertson C. Measuring the effect of enhanced cleaning in a UK hospital: a prospective cross-over study. *BMC Medicine* 2009;7:28–40.
- Boyce JM, Havill NL, Lipka A, Havill H, Rizvani BS. Variations in hospital daily cleaning practices. *Infect Control Hosp Epidemiol* 2010;31:99–101.
- Hota B, Blom DW, Lyle EA, Weinstein RA, Hayden MK. Interventional evaluation of environmental contamination by vancomycin-resistant enterococci: failure of personnel, product, or procedure? *J Hosp Infect* 2009;71:123–131.
- Mulvey D, Redding P, Robertson C, et al. Finding a benchmark for monitoring hospital cleanliness. *J Hosp Infect* 2011;77:25–30.
- Carling PC, Briggs JL, Perkins J, Highlander D. Improved cleaning of patient rooms using a new targeting method. *Clin Infect Dis* 2006;42:385–388.
- Rutala WA, Weber DJ; the Healthcare Infection Control Practices Advisory Committee (HICPAC), Centers for Disease Control and Prevention. Guideline for disinfection and sterilization in healthcare facilities, 2008. http://www.cdc.gov/hicpac/pdf/guidelines/Disinfection_Nov_2008.pdf. Accessed November 15, 2012.
- Sherlock O, O'Connell N, Creamer E, Humphreys H. Is it really clean? an evaluation of the efficacy of four methods for determining hospital cleanliness. *J Hosp Infect* 2009;72:140–146.
- Carling PC, Parry MM, Rupp ME, Po JL, Dick B, Von Behren S. Improving cleaning of the environment surrounding patients in 36 acute care hospitals. *Infect Control Hosp Epidemiol* 2008; 29:1035–1041.

Improving Hand Hygiene Compliance in Hospitals by Design

Essential in reducing hospital-acquired infections is adequate hand hygiene (HH) among healthcare workers (HCWs).¹ International studies show, however, that HH guidelines are adhered to in less than 50% of required times.² Research into HH behavior has shown that self-reported compliance is often higher than observed compliance, which seems to indicate that HCWs are unaware of their HH behavior.² In addition, because of its frequency, HH behavior could be considered an automatic (or subconscious) behavior.³ Therefore, a (temporary) shift from the subconscious to the conscious could be a solution to change current HH behavior and create new habits. These insights formed the points of departure of the design project described here, which aimed to develop an alcohol-based hand rub (ABHR) dispenser to stimulate

HCWs to better adhere to the international guidelines of HH.^{4,5} In order to increase the chance of success of the new dispenser, a participatory design approach was applied, meaning that all stakeholders of the dispenser (nurses, physicians, infection control practitioners, housekeeping) were actively involved in the different phases of the development process.^{6,7}

The development process of the new dispenser consisted of 5 phases: analysis, idea finding (identifying promising design directions), and 3 idea iterations. Interim ideas were evaluated using functional 3-dimensional prototypes and the results applied to further develop the final concept. Methods of user research included observations and individual and focus group interviews.

The new ABHR dispenser that resulted from this project incorporates 3 distinctive features: (1) instant feedback, (2) a spray mechanism, and (3) an integrated drip tray (see Figure 1). First, the new dispenser provides instant feedback to the HCWs on their frequency of using the dispenser in order to increase their awareness about their HH behavior. By means of 5 light-emitting diodes (LEDs), the number of performed HH moments is displayed during a fixed period of time (15 minutes). The LED feedback refers symbolically to the 5 moments of HH as prescribed by the World Health Organization.⁸ This feedback, which does not differentiate between multiple users, is expected to stimulate a higher frequency of use by acting as a mirror (see what your behavior is) and a mediator (see the behavior of your colleagues and discuss it). When more than 5 device activations occur within the specified 15 minutes, the LED circle starts again. Second, the dispenser uses a spray mechanism that sprays a fixed amount

(3 mL) of ABHR on the back of both hands. This way, alcohol is applied directly between the fingers, an important part of the hand that is difficult to clean.⁵ The dispenser contains 2 refills to ensure that the device is less likely to run empty. Third, the integrated drip tray allows for the possibility of universal placement (ie, not restricted to the sink area) without damage to floors or other surfaces.

Twenty nurses and 4 physicians working in the intensive care unit and surgical ward of a large university teaching hospital in the Netherlands tested the new dispenser. The test materials consisted of a functional prototype (Figure 1, right) and an evaluation questionnaire covering topics on access, performance feedback, stimulation of use, social control, professional look and feel of the product, and ranking of the 3 distinctive design features. The test took place in a 4-, 2-, or 1-patient room. HCWs were asked to use the prototype and report their experiences to the researcher and to complete the questionnaire. All user tests were audio-recorded and summarized. Questionnaires were analyzed in SPSS (v15). Each item was scored on a 7-point Likert scale, except for the most important feature of the dispenser, which was scored by ranking.

Most participants considered the spray functionality easy to use and appreciated that the correct amount of ABHR was dispensed automatically. The participants mentioned that the spray mechanism made the product appearance more luxurious. Sometimes it was ambiguous for users how their hands should be placed in the dispenser (palms up or down). Participants appreciated the way the concept provided performance feedback and considered it useful in combination

FIGURE 1. Visualization of the final concept of the newly developed alcohol dispenser (left) and the functional prototype used for user testing (right).⁵ Distinctive features of the dispenser are (1) instant feedback, (2) spray mechanism, and (3) integrated drip tray. A color version of this figure is available in the online edition of *Infection Control and Hospital Epidemiology*.

with education. However, they were uncertain whether it would continue to hold attention over time.

The questionnaire confirmed that participants were positive toward the design. In particular, participants reported that the dispenser was a positive addition to the patient room (median, 5; interquartile range [IQR], 1.5), with a professional look (median, 5; IQR, 2) enabling easy access and use (median, 5; IQR, 2). Participants reported that the dispenser gave insight into their performance and helped them adhere to the protocol (median, 5; IQR, 2), although they were neutral as to whether the dispenser would help colleagues address each other on their behavior (median, 4; IQR, 3). The participants considered the spray functionality the most important feature of the concept (first place, 46%), closely followed by the integrated drip tray (first place, 42%). The LED performance feedback was ranked as the least important feature.

The study was limited by the scope, namely, a 1-center study, with data collection having to be fitted into already busy shifts. User tests were therefore always kept as short as possible, and only 4 physicians could be included in the sample. On the other hand, participation of HCWs in the study activities has yielded unique insights into workable solutions, which have been incorporated in the design. Currently, a follow-up study is being conducted on the optimal location of alcohol dispensers within patient rooms.

ACKNOWLEDGMENTS

Potential conflicts of interest. All authors report no conflicts of interest relevant to this article. All authors submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest, and the conflicts that the editors consider relevant to this article are disclosed here.

Marijke Melles, PhD;¹ Vicki Erasmus, PhD;²
Martijn P. M. van Loon, MSc;¹ Marc Tassoul, MSc;¹
Ed F. van Beeck, MD, PhD;² Margreet C. Vos, MD, PhD³

Affiliations: 1. Delft University of Technology, Faculty of Industrial Design Engineering, Delft, The Netherlands; 2. Erasmus University Medical Centre, Department of Public Health, Rotterdam, The Netherlands; 3. Erasmus University Medical Centre, Department of Medical Microbiology and Infectious Diseases, Rotterdam, The Netherlands.

Address correspondence to Marijke Melles, PhD, Landbergstraat 15, 2628 CE Delft, The Netherlands (m.melles@tudelft.nl).

Received February 1, 2012; accepted August 9, 2012; electronically published November 19, 2012.

Presented in part: Proceedings of the International Conference on Healthcare Systems, Ergonomics and Patient Safety; Oviedo, Spain; June 2011.

Infect Control Hosp Epidemiol 2013;34(1):102–104

© 2012 by The Society for Healthcare Epidemiology of America. All rights reserved. 0899-823X/2013/3401-0018\$15.00. DOI: 10.1086/668772

REFERENCES

1. Donaldson L. *Dirty Hands: The Human Cost*. London: UK Department of Public Health, 2006.
2. Erasmus V, Dahan TJ, Brug J, et al. A systematic review of studies on compliance to hand hygiene guidelines in health care. *Infect Control Hosp Epidemiol* 2010;31(3):283–294.
3. Verplanken B, Aarts H. Habit, attitude and planned behaviour: is habit an empty construct or an interesting case of goal-directed automaticity? *Eur Rev Soc Psychol* 1999;101–134.
4. Allegranzi B, Storr J, Dziekan G, Leotsakos A, Donaldson L, Pittet D. The first global patient safety challenge “clean care is safer care”: from launch to current progress and achievements. *J Hosp Infect* 2007;65(suppl 2):115–123.
5. Van Loon M. *Getting Hands Clean: Developing a Handhygiene Reminder for the Intensive Care Unit and the Surgical Ward*. Delft, Netherlands: Delft University of Technology, 2009.
6. Schuler D, Namioka A. *Participatory Design: Principles and Practices*. Hillsdale, NJ: Lawrence Erlbaum, 1993.
7. Preece J, Rogers Y, Sharp H. *Interaction Design: Beyond Human-Computer Interaction*. New York: Wiley, 2002.
8. Sax H, Allegranzi B, Uckay I, Larson E, Boyce J, Pittet D. “My five moments for hand hygiene”: a user-centred design approach to understand, train, monitor and report hand hygiene. *J Hosp Infect* 2007;67(1):9–21.