Embedding silicon spin qubits in superconducting circuits

Zheng, G.; Samkharadze, N.; Noordam, M. L.; Kalhor, N.; Brousse, D.; Sammak, A.; Mendes, U. C.; Blais, A.; Scappucci, G.; Vandersypen, L. M.K.

DOI

[10.1364/QIM.2019.F3B.3](https://doi.org/10.1364/QIM.2019.F3B.3)

Publication date

2019

Document Version

Final published version

Published in

Quantum Information and Measurement, QIM 2019

Citation (APA)

Zheng, G., Samkharadze, N., Noordam, M. L., Kalhor, N., Brousse, D., Sammak, A., Mendes, U. C., Blais, A., Scappucci, G., & Vandersypen, L. M. K. (2019). Embedding silicon spin qubits in superconducting circuits. In *Quantum Information and Measurement, QIM 2019* [142378] (Optics InfoBase Conference Papers; Vol. Part F165-QIM 2019). OSA - The Optical Society. <https://doi.org/10.1364/QIM.2019.F3B.3>

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Green Open Access added to TU Delft Institutional Repository

'You share, we take care!' - Taverne project

<https://www.openaccess.nl/en/you-share-we-take-care>

Otherwise as indicated in the copyright section: the publisher is the copyright holder of this work and the author uses the Dutch legislation to make this work public.

Embedding Silicon Spin Qubits in Superconducting Circuits

G. Zheng^{1,*}, N. Samkharadze¹, M. L. Noordam¹, N. Kalhor¹, D. Brousse², A. Sammak², U. C. Mendes³, A. Blais^{3,4}, G. Scappucci¹, L. M. K. Vandersypen^{1,†}

¹*QuTech and Kavli Institute of Nanoscience, Delft University of Technology Lorentzweg 1, 2628 CJ Delft, Netherlands.*

²*QuTech and Netherlands Organization for Applied Scientific Research (TNO), Stieltjesweg 1 2628 CK Delft, Netherlands.*

³*Institut Quantique and Département de Physique, Université de Sherbrooke, Sherbrooke, Québec J1K 2R1, Canada.*

⁴*Canadian Institute for Advanced Research, Toronto, Ontario, Canada.*

*g.zheng@tudelft.nl, †l.m.k.vandersypen@tudelft.nl

Abstract: We demonstrate the strong coupling between a single electron spin in silicon and a single photon in a superconducting microwave cavity. Using the same cavity we perform rapid high-fidelity single-shot readout of two-electron spin states. © 2019 The Author(s)

OCIS codes: 270.5585, 270.5580.

Long coherence times of single spins in silicon quantum dots make these systems highly attractive for quantum computation. However, the realization of large networks of quantum dot-based spin qubits requires advancements in coherent long-range interconnects between the qubits, and more efficient use of the on-chip real estate [1]. A promising approach that addresses these challenges is to integrate spin qubits in a superconducting cavity.

While direct interactions between quantum dot-based spin qubits are limited to a submicron range, circuit quantum electrodynamics can be employed to extend this range to millimeter scale, by transferring quantum information via microwave photons. Here, we report the strong coupling between a single electron spin in a silicon double quantum dot and a microwave photon in an on-chip high-impedance superconducting cavity [2–5], a crucial requirement for the coherent cavity-mediated transfer of quantum information between distant spin qubits. The electric field component of the cavity couples directly to the dipole moment of the electron charge in the double quantum dot, and indirectly to the electron spin, through a local magnetic field gradient from a nearby micromagnet [6, 7]. A charge-photon coupling strength of 200 MHz and a spin-photon coupling strength as large as 15 MHz have been extracted, several times larger than the spin and cavity decay rates.

Another crucial requirement for operating spin-based quantum processors is the ability to read out spin qubits fast and in a single shot manner. So far, this has been achieved using nearby single-electron transistors (SETs) as charge detectors, in combination with a spin-to-charge conversion scheme [8]. However, these SETs come with additional resources (gate electrodes, electron reservoirs), which poses an additional hurdle for scaling up to dense spin qubit systems. An alternative approach is to connect a resonant circuit to gates that are already in place for confining electrons [9–11]. Here, we utilize the cavity to detect the charge susceptibility at a two-electron interdot transition. Using Pauli's exclusion principle to map charge to spin information, we demonstrate the gate-based readout of a two-electron spin state in a single shot with an average fidelity of >98% in 6 μ s.

These results open the way to coherent long-range interactions between spin qubits and rapid high-fidelity readout using the same cavity, removing a major roadblock for the scalability of spin-based quantum processors.

1. References

References

1. L. M. K. Vandersypen, H. Bluhm, J. S. Clarke, A. S. Dzurak, R. Ishihara, A. Morello, D. J. Reilly, L. R. Schreiber, and M. Veldhorst, *npj Quantum Inf.* 3, 34 (2017).
2. N. Samkharadze, A. Bruno, P. Scarlino, G. Zheng, D.P. DiVincenzo, L. DiCarlo, and L.M.K. Vandersypen, *Phys. Rev. Applied* 5, 044004 (2016).
3. N. Samkharadze, G. Zheng, N. Kalhor, D. Brousse, A. Sammak, U. C. Mendes, A. Blais, G. Scappucci, L. M. K. Vandersypen, *Science* 359, 6380 (2018).
4. X. Mi, M. Benito, S. Putz, D. M. Zajac, J. M. Taylor, Guido Burkard, and J. R. Petta, *Nature* 555, 599-603 (2018).

5. A. J. Landig, J. V. Koski, P. Scarlino, U. C. Mendes, A. Blais, C. Reichl, W. Wegscheider, A. Wallraff, K. Ensslin, and T. Ihn, *Nature* 560, 179184 (2018).
6. X. Hu, Y.-X. Liu, and F. Nori, *Phys. Rev. B* 86, 035314 (2012).
7. F. Beaudoin, D. Lachance-Quirion, W. A. Coish, M. Pioro-Ladrère, *Nanotechnology* 27, 464003 (2016).
8. R. Hanson, L. P. Kouwenhoven, J. R. Petta, S. Tarucha, and L. M. K. Vandersypen, *Rev. Mod. Phys.* 79, 1217 (2007).
9. A. West, B. Hensen, A. Jouan, T. Tanttu, C.H. Yang, A. Rossi, M.F. Gonzalez-Zalba, F.E. Hudson, A. Morello, D.J. Reilly, A.S. Dzurak, arXiv:1809.01864 (2018).
10. P. Pakkiam, A. V. Timofeev, M.G. House, M.R. Hogg, T. Kobayashi, M. Koch, S. Rogge, M.Y. Simmons, arXiv:1809.01802 (2018).
11. M. Urdampilleta, D. J. Niegemann, E. Chanrion, B. Jadot, C. Spence, P.-A. Mortemousque, C. Buerle, L. Hutin, B. Bertrand, S. Barraud, R. Maurand, M. Sanquer, X. Jehl, S. De Franceschi, M. Vinet, T. Meunier, arXiv:1809.04584 (2018).