

The discharge magnitude of the 1374 millennium flood event in the Rhine river

Ngo, Hieu; Bomers, Anouk; Augustijn, Dennie CM.; Ranasinghe, Roshanka; Filatova, T.; Hulscher, Suzanne J.M.H.

Publication date

2022

Document Version

Final published version

Published in

Anthropogenic Rivers: Book of Abstracts NCR DAYS 2022 13-14 April | TU Delft

Citation (APA)

Ngo, H., Bomers, A., Augustijn, D. CM., Ranasinghe, R., Filatova, T., & Hulscher, S. J. M. H. (2022). The discharge magnitude of the 1374 millennium flood event in the Rhine river. In A. Blom, L. M. Stancanelli, J. A. Dercksen, C. Ylla Arbós, M. K. Chowdhury, S. M. Ahrendt, C. Piccoli, R. M. J. Schielen, K. Sloff, & J. H. Slinger (Eds.), *Anthropogenic Rivers: Book of Abstracts NCR DAYS 2022 13-14 April | TU Delft* (pp. 14-15). (NCR Publication; No. 49-2022).

Important note

To cite this publication, please use the final published version (if applicable).
Please check the document version above.

Copyright

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

Takedown policy

Please contact us and provide details if you believe this document breaches copyrights.
We will remove access to the work immediately and investigate your claim.

Anthropogenic Rivers

Book of Abstracts

NCR DAYS 2022

13-14 April | TU Delft



*Astrid Blom, Laura M. Stancanelli, Jelle A. Dercksen, Clàudia Ylla Arbós,
M. Kifayath Chowdhury, Shelby M. Ahrendt, Carolina Piccoli,
Ralph M.J. Schielen, Kees Sloff & Jill H. Slinger (eds.)*

NCR Publication: 49-2022

Netherlands
Centre for
River studies **NCR**

The discharge magnitude of the 1374 millennium flood event in the Rhine river

Hieu Ngo^{a,b}, Anouk Bomers^b, Dennie CM. Augustijn^b, Roshanka Ranasinghe^{b,c}, Tatiana Filatova^d, and Suzanne JMH. Hulscher^b

^a University of Twente, Department of Governance and Technology for Sustainability, Drienerloaan 5, 7522 NB, Enschede, the Netherlands

^b University of Twente, Department of Water Engineering and Management, Drienerloaan 5, 7522 NB, Enschede, the Netherlands

^c IHE Delft, Coastal & Urban Risk & Resilience department, Westvest 7, 2611 AX, Delft, the Netherlands

^d Delft University of Technology, Multi Actor Systems Department, Delft, 2628 CN, the Netherlands

Keywords — historic flood event, hydraulic modelling approach, flood reconstruction, flood frequency analysis

Introduction

Reconstructions of the most severe historic flood events contribute to a better quantification of design discharges corresponding to large return periods. However, reconstructions of the peak discharges of these historic flood events generally have large uncertainties related to the historic topography and hydraulic roughness of the river channels and floodplains (Herget and Meurs, 2010; Lang et al., 2003).

Discharge magnitudes of historic flood events can be reconstructed by using the simple one-dimensional (1D) approach (e.g. 1D cross-sectional) (Herget and Meurs, 2010) or hydraulic modelling approach (van der Meulen et al., 2021). However, the simple 1D approach makes it difficult to account for uncertainties in historic topography and hydraulic roughness. Therefore, this study sets up a 1D-2D coupled hydraulic model to reconstruct the discharge magnitude of the 1374 flood event - considered the largest flood of the last millennium in the Rhine river - in which the river is modelled by 1D profiles and the floodplains are discretized on a 2D grid. The result will be used to evaluate the effect on the design discharge.

Methodology

An “inverse modelling” approach was used to reconstruct the discharge magnitude of the 1374 flood event. First, the 1D-2D coupled hydraulic model was set up with the historic topography of the main river and floodplains corresponding to the year 1374 extracted from a high-resolution palaeo-DEM for the Lower Rhine catchment for 800 AD (van der Meulen et al., 2020) (Fig 1) and the hydraulic roughness coefficient for land cover classes corresponding to the palaeo situation (van der Meulen et al., 2021). An uncertainty analysis was then

performed with different river bed levels and hydraulic roughness values (Table 1) to estimate the influence of these uncertainties on the reconstructed peak discharge of the 1374 flood event. The upstream discharge wave was also varied, corresponding to a wide range of peak discharges (12,000-24,000 m³/s). Next, the simulated water levels were compared with the flood mark (observed water level) corresponding to the 1374 flood event at Cologne to determine the appropriate discharge magnitude.




Figure 1. The model set-up for the study area (left side) and a close-up of the 2D grid (right side).

Table 1. Landscape classes and assigned Manning's n values. Source: van der Meulen et al. (2021)

Class	n_{\min}	n_{average}	n_{\max}
High grounds	0.1	0.1	0.1
River bed and banks	0.025	0.03	0.045
Proximal floodplain	0.06	0.07	0.08
Distal floodplain, high	0.04	0.05	0.06
Distal floodplain, low	0.035	0.04	0.055

Results and Discussion

The upstream discharge magnitude of the 1374 flood event was determined to be between 12,500 m³/s and 22,000 m³/s considering the combinations of the maximum and minimum roughness coefficient values with different river bed levels, with the ‘best’ estimate between 14,600 m³/s and 18,700 m³/s corresponding to

* Corresponding author
Email address: h.q.ngo@utwente.nl (Hieu Ngo)

the average roughness coefficient value (n_{average}) for all landscape classes.




Figure 2. Water level (WL) at Cologne plotted against peak discharge. The uncertainty bands at peak discharge between 12,000 and 24,000 m³/s show results with all roughness classes set to minimum and maximum Manning's n values corresponding to river bed level = 800 AD-3 m. The red marker shows the WLs corresponding to the average Manning's n values.

The reconstructed discharge magnitude of the 1374 flood event corresponds to the study of Herget and Meurs (2010) (23,200 m³/s, with an uncertainty range of between 18,800 m³/s and 29,000 m³/s) and this study were independently updated into the 12 reconstructed historic flood events in the period 1300-1772. Then these data were respectively combined with the systematic discharge data set covering the period 1772-2018 to create a continuous discharge data set covering the period 1317-2018 corresponding to each study by using a bootstrap method (Bomers et al., 2019). A flood frequency analysis was performed based on these discharge data sets to determine the design discharges and their 95% confidence interval for different return periods corresponding to each study based on the generalized extreme value (GEV) distribution function (Fig 3).

The results (Fig 3) show that updating the 1374 reconstructed discharge magnitude of this study

into the historic flood events results in a significant reduction of 1,800 m³/s (10%) in the design discharge and 1000 m³/s (20%) in the confidence interval corresponding to a 100,000 year return period compared to using the 1374 reconstructed discharge magnitude of Herget and Meurs (2010).

Conclusions

The discharge magnitude for the 1374 millennium flood event was determined to be between 12,500 m³/s and 22,000 m³/s, with the 'best' estimate between 14,600 m³/s and 18,700 m³/s.

The reduction of the 95% confidence interval shows the importance of reconstructing historic flood events with high accuracy, contributing a more certain quantification of design discharges.

References

- Bomers, A., Schielen, R.M.J., Hulscher, S.J.M.H. (2019) Decreasing uncertainty in flood frequency analyses by including historic flood events in an efficient bootstrap approach. *Nat Hazards Earth Syst Sci* 19:1895–1908. <https://doi.org/10.5194/nhess-19-1895-2019>
- Herget, J., Meurs, H. (2010) Reconstructing peak discharges for historic flood levels in the city of Cologne, Germany. *Global and Planetary Change* 70:108–116. <https://doi.org/10.1016/j.gloplacha.2009.11.011>
- Lang, M., Moussay, D., Recking, A., Naulet, R. (2003) Hydraulic modelling of historical floods. Case study on the Ardeche river at Vallon-Pont-d Arc. In Thorndycraft, V.R., Benito, G., Barriendos, M., Llasat, C. (eds.) *Palaeofloods, Historical Data & Climatic Variability: Applications in Flood Risk Assessment* (Proceedings of the PHEFRA International Workshop, Barcelona, 16-19th. October 2002) CSIC Madrid pp.183-189
- van der Meulen, B., Cohen, K.M., Pierik, H.J, et al (2020) LiDAR-derived high-resolution palaeo-DEM construction workflow and application to the early medieval Lower Rhine valley and upper delta. *Geomorphology* 370:107370. <https://doi.org/10.1016/j.geomorph.2020.107370>
- van der Meulen, B., Bomers, A, Cohen, K.M., Middelkoop, H. (2021) Late Holocene flood magnitudes in the Lower Rhine river valley and upper delta resolved by a two-dimensional hydraulic modelling approach. *Earth Surface Processes and Landforms* 46:853–868. <https://doi.org/10.1002/esp.5071>





Fig 3. Flood frequency curves and their 95% confidence intervals of data sets corresponding to the reconstructed discharge magnitude of the 1374 flood event of Herget and Meurs's study and this study.