

## **Celebración de 50 años de SWIM (Salt Water Intrusion Meetings – Reuniones sobre la Intrusión de Agua Salada)**

Post, Vincent E.A.; Essink, Gualbert Oude; Szymkiewicz, Adam; Bakker, Mark; Houben, Georg; Custodio, Emilio; Voss, Clifford

**DOI**

[10.1007/s10040-018-1800-8](https://doi.org/10.1007/s10040-018-1800-8)

**Publication date**

2018

**Document Version**

Accepted author manuscript

**Published in**

Hydrogeology Journal

**Citation (APA)**

Post, V. E. A., Essink, G. O., Szymkiewicz, A., Bakker, M., Houben, G., Custodio, E., & Voss, C. (2018). Celebración de 50 años de SWIM (Salt Water Intrusion Meetings – Reuniones sobre la Intrusión de Agua Salada). *Hydrogeology Journal*, 26(6), 1767–1770 . <https://doi.org/10.1007/s10040-018-1800-8>

**Important note**

To cite this publication, please use the final published version (if applicable).  
Please check the document version above.

**Copyright**

Other than for strictly personal use, it is not permitted to download, forward or distribute the text or part of it, without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license such as Creative Commons.

**Takedown policy**

Please contact us and provide details if you believe this document breaches copyrights.  
We will remove access to the work immediately and investigate your claim.

# Celebrating 50 years of SWIMs (Salt Water Intrusion Meetings)

*Vincent E. A. Post<sup>1,2,\*</sup>, Gualbert Oude Essink<sup>3,4</sup>, Adam Szymkiewicz<sup>5</sup>, Mark Bakker<sup>6</sup>, Georg Houben<sup>1</sup>, Emilio Custodio<sup>7</sup>, Clifford Voss<sup>8</sup>*

*<sup>1</sup>Federal Institute for Geosciences and Natural Resources (BGR), Hannover, Germany*

*<sup>2</sup>College of Science and Engineering, Flinders University, Adelaide, Australia*

*<sup>3</sup>Deltares, Utrecht, the Netherlands*

*<sup>4</sup>Department of Physical Geography Utrecht University, Utrecht, the Netherlands*

*<sup>5</sup>Department of Geotechnics, Geology and Marine Civil Engineering, Faculty of Civil and Environmental Engineering, Gdańsk University of Technology, Gdańsk, Poland*

*<sup>6</sup>Faculty of Civil Engineering and Geosciences, TU Delft, Delft, the Netherlands*

*<sup>7</sup> Department of Civil and Environmental Engineering. Technical University of Catalonia (UPC), Barcelona, Spain*

*<sup>8</sup>U.S. Geological Survey, Menlo Park, California, USA*

*\*Contact author e-mail address: [vincent.post@bgr.de](mailto:vincent.post@bgr.de)*

## Keywords

salt-water/fresh-water relations, salt-water intrusion, salinization, history of hydrogeology

**NOTE TO COPYEDITOR – PLEASE INSERT THE FOLLOWING AS A FIRST-PAGE FOOTNOTE:**

This article is part of the topical collection “Celebrating 50 years of SWIMs (Salt Water Intrusion Meetings)”

## Abstract

The Salt Water Intrusion Meetings, or SWIMs, are a series of meetings that focus on seawater intrusion in coastal aquifers and other salinization processes. 2018 marks the 50<sup>th</sup> year of the SWIM and the 25<sup>th</sup> biennial meeting. The SWIM proceedings record half a century of research progress on site characterisation, geophysical and geochemical techniques, variable-density flow, modelling, and water management. The SWIM is positioning itself to remain a viable platform for coastal aquifer management challenges of the next 50 years.

## **1. Emergence of SWIM**

Almost fifty years ago, on 28 and 29 November in 1968, a group of 20 German, Danish and Dutch scientists met in Hannover, Germany, to discuss salinization of groundwater in coastal areas. They had been invited by the late Prof. Wolfgang Richter (working at the Geological Survey of Lower Saxony), who recognised the need to exchange knowledge with neighbouring countries that faced similar saline groundwater problems. Participant contributions were printed and most were gathered in a proceedings report. Due to the usefulness of the meeting, another meeting was organized in the Netherlands. The conference series, now known as the Salt Water Intrusion Meeting, was born.

The second meeting (13 to 15 May 1970 in Vogelenzang, the Netherlands) was organised by the Dutch committee of UNESCO's International Hydrological Decade (IHD, 1965 to 1974). This meeting had a logo, the outline of the organising country visible through the wavy lines of the UNESCO IHD logo (Figure S1 of the electronic supplementary material (ESM)). The name 'Salt Water Intrusion Meeting' was introduced at the third meeting (1972 in Copenhagen, Denmark). The conference series was maturing with a catchy acronym (SWIM) and a logo style that has been used for almost all following SWIMs.

## **2. Topics and trends**

The first proceedings describes the early application of techniques that have become commonplace today. While most contributions focussed on describing the fresh-saline groundwater situation using hydrochemical data, contributions about geo-electrical techniques were also prominently represented. Computers were already used to make resistivity models with more than three layers (Flathe, 1968). Quite sophisticated chemical tracer techniques were being applied, and the first proceedings contained contributions on the use of isotopes (Stahl, 1968, Nielsen, 1968, Geyh, 1968) to infer origins of different coastal groundwater types.

The current pervasiveness of numerical modelling and software tools was still years away. The first SWIM contribution about using numerical techniques was by Verruijt (1972), who demonstrated the principles of the finite element method and showed two applications to theoretical freshwater-

saltwater interface problems. Presentations about numerical models of real-world coastal aquifers took another decade to appear.

Case studies describing coastal aquifer systems have been part of SWIMs since the first meeting and constitute an important contribution to the knowledge of coastal areas. As the number of participating countries grew, the geographical diversity of contributions increased (Figure 1). At the first two meetings, the focus was on unconsolidated sedimentary aquifers prevalent in the three organising countries. The first descriptions of salinization in chalk aquifers soon followed (e.g. Downing, 1972). Karst systems were extensively discussed at the Hannover meeting in 1979, the proceedings of which feature a paper of 68 pages about the coastal aquifer system of Apulia, Italy (Cotecchia, 1981). Several case studies in fractured bedrock aquifers were presented at the 1981 Sweden meeting (e.g. Lindewald, 1981; Sund and Bergman, 1981). A few SWIM contributions deal with the movement of saline groundwater in inland aquifers, for example, the genesis and migration of brines originating from leaching of salt domes or stratiform salt deposits (e.g. Ochmann and Fielitz 1993; Klinge et al. 1999; Zechner et al. 2014), and the existence of brackish groundwater due to aridity (Custodio, 1992).

In a landmark early SWIM paper, Lebbe (1981) demonstrated the existence of an intruded seawater cell below a wide beach in Belgium surrounded in the subsurface by fresher groundwater. He recognised that the circulatory flow of intruded seawater inside the cell was driven by the tide and used a numerical model to quantitatively interpret flow dynamics. Since then, similar tidally-driven circulation cells have been found elsewhere, but Lebbe's (1981) SWIM contribution can be credited as the first description of such a system. More generally, interactions between coastal aquifers and the ocean and submarine groundwater discharge have become a main SWIM theme. Early contributions dealt mainly with submarine springs in karst systems (Schwerdtfeger, 1981), while more recently interest has shifted towards the role of submarine groundwater discharge as an important source of nutrients to the marine environment (e.g. Scholten et al., 2014).

Some interesting trends are visible when the contributions to the meetings are aggregated in categories (Figure S2 of the ESM). The mathematical treatment of sharp interface solutions, frequently discussed at early meetings, has become a rare topic. At the first two meetings, geochemistry comprised a considerable proportion of the contributions, then disappeared during the 1970s, and increased again, reaching a peak (as a percentage) in the mid 1990s. Geophysics, (numerical) modelling, management, sea-level rise, island hydrology, and climate change, have become more prominent topics, although the latter category has been showing a decline since 2010.

### **3. From European to global**

The 1960s world was not yet as globalised as today, so SWIM began as a west-European meeting series and contributions by non-European participants remained rare during the meetings organised before 2000. At the 1986 SWIM in Delft, the Netherlands, Cliff Voss (US Geological Survey) suggested to make SWIM a worldwide conference because of its great importance to all countries, but the European organizers did not wish to make it more difficult for European students, who often gave their first international talk at the SWIMs, to travel to the meetings. This later led to the establishment of the Salt Water Intrusion and Coastal Aquifer (SWICA) conference series that were held in Essaouira, Morocco, in 2001 and in Merida, Mexico, in 2003. Since both SWIM and SWICA meetings attracted the same participants, it was discussed during the 18<sup>th</sup> SWIM that the biennial meetings be combined, with meeting location alternating between a European country (just as in the original SWIM series) and a non-European country. The first instance of the new combination was in 2006, a joint 19<sup>th</sup> SWIM and the 3<sup>rd</sup> SWICA meeting held in Cagliari, Italy, at which the attendees voted to combine the two meeting series under the name SWIM.

Since 2009, Asian countries have been organising a successful series, the Asia-Pacific Coastal Aquifer Management Meeting (APCAMM). The first APCAMM was held in Bangkok, Thailand. The 24<sup>th</sup> SWIM and the 4<sup>th</sup> APCAMM were held jointly in Cairns, Australia, in 2016. There are no current plans to merge with SWIM and APCAMM, as APCAMM has a specific geographic focus, but there will be coordination, joint participation, and exchange of ideas between these conference series.

## **4. Current status**

SWIM has become a conference series with a strong reputation. The format has evolved into a one-week meeting, always with exclusively plenary sessions, and a mid-week field trip to sites related to saltwater problems. In recent years it has become customary to organise short courses, for example on seawater intrusion modelling, prior to the meeting. At the 24<sup>th</sup> SWIM in Cairns, the participants voted to continue the plenary-only format for all future meetings, as this exchange of ideas among all attendees is one of the attributes that makes SWIM so successful.

The informal passing of the responsibility of hosting the next meeting to the next organizer has proven to be a sustainable model that has ensured the SWIM continuity for half a century. The number of participants has risen over the years (peaking at 165 in Husum, Germany, in 2014, see Figure S3 of the ESM), while the SWIM's primary objective is still the same as the first meeting: the exchange of knowledge on saline groundwater problems.

The 25<sup>th</sup> SWIM in Gdańsk, Poland, from 17 to 22 June 2018 (see <http://www.swim2018.syskonf.pl>) is the fifth SWIM organized in a Baltic country. Despite the low salinity of the Baltic Sea relative to ocean water, numerous cases of saltwater intrusion have been investigated in this region, from spits

(e.g. Dowgiałło & Frączek 1990) to deltaic areas (e.g. Zuber et al. 1990) to crystalline bedrock islands (e.g. Lindewald 1981). A previous SWIM meeting in Sopot, a locality near Gdańsk, took place in 1990. Great socio-economic changes began at that time in Eastern Europe, which, among many other consequences, led to a significant shift in groundwater usage patterns in the Gdańsk region and Poland. The risk of salinization of coastal water wells substantially decreased due to reduced water uptake by heavy industry while contamination by nutrients originating from agricultural practices has become a major concern (e.g. Højberg et al. 2017, Szymczycha & Pempkowiak 2016).

The proceedings of all SWIMs are available online and can be accessed via the website <http://www.swim-site.org> (SWIM 2018). They archive developments in coastal hydrogeology characterization, process understanding, and water-supply management over the past half century. The causes of salinization are now better understood than at the first meeting, and the technologies to measure and model groundwater processes in coastal areas, albeit still in need of improvement, have reached high levels of sophistication. A special volume republishing selected papers from the SWIMs prior to 1990 was published by the International Association of Hydrogeologists (De Breuck, 1991). A table with a listing of all SWIM and SWICA conferences is given in Table S1 of the ESM.

## 5. The next 50 years

The problems of saltwater intrusion and aquifer salinization have not been solved. Population growth, urbanisation, land-use change, environmental destruction, globalisation, and tourism are increasing the pressure on coastal freshwater resources (Michael et al. 2017). The possible acceleration of sea-level rise and land subsidence compound the seawater intrusion problems in coastal areas. At the same time, monitoring networks are often inadequate, leading to a paucity of data for numerical model analyses and management decisions. Governance issues such as a lack of adequate legislation and enforcement, as well as poor users' and civil society involvement form other obstacles against effectively combating saltwater intrusion problems, let alone the more general problem of the slowness of the uptake of new science into management and society.

There are many important topics for which better understanding of the physical system is required, as evidenced by recent trends in SWIM contributions. The example of submarine groundwater discharge was already mentioned in the context of the Baltic Sea, but to understand its importance to ocean water quality and ecosystems across the globe is an essential current research challenge. The topic of land-aquifer-ocean connection is likely to see significant advances in the future. Research into better ways of describing mixing in groundwater, and the effects of heterogeneity (as well as tides and wave action) on coastal aquifer water quality, also continues as contributions to SWIM. All these new physical insights, partly based on increasingly sophisticated measurement techniques, will require

more sophisticated numerical models that could certainly benefit from quantum computers when they become reality.

The relevance of the SWIM is thus greater than ever. One future challenge may be to partly transform the meeting from a platform where experts discuss problems to one where solutions are explored and developed. An increasing number of SWIM contributions is shifting towards providing solutions, such as managed aquifer recharge, the control of saline groundwater, the capture of marine water through the aquifer to feed desalination plants, and the use of coastal aquifers for the storage of surplus water. Bridging the gap between science and management is another direction that SWIM will likely take, and in this perspective, SWIMs will likely develop more outreach activities. The recent video that was produced by GEUS in Denmark is a good example in that sense (accessible via SWIM 2018). So, apart from celebrating SWIM's 50th anniversary, the 2018 meeting in Gdańsk will also be used to develop ideas about how SWIM can continue its successful tradition and at the same time evolve and innovate to continue its relevance in the coming five decades, which, however scary it may sound, may see even faster and more far-reaching global changes than the past 50 years.

## References

- Bauer, P., S. Zimmermann, R. Held, T. Gumbrecht and W. Kinzelbach (2002) Is density flow balancing the salt budget of the Okavango delta? Evidence from field and modelling studies. SWIM 17, 309 – 321.
- Cotecchia, V. (1981) Methodologies adopted and results achieved in the investigations of seawater intrusion into the aquifer of Apulia (Southern Italy). *Geologisches Jahrbuch Reihe C* 29 (SWIM6), 1 – 68.
- Custodio, E (1992) Coastal aquifer salinization as a consequence of aridity: the case of Amurga phonolitic massif, Gran Canaria island. SWIM 12, 81 – 98.
- De Breuck, W. [editor] (1991). *Hydrogeology of Salt Water Intrusion: A Selection of SWIM Papers, International Contributions to Hydrogeology, Volume 11*, International Association of Hydrogeologists, Verlag Heinz Heise, 422 p.
- Dowgiałło J., Frączek E. (1990) An attempt at the interpretation of new data on the Hel Spit hydrogeology (Poland). SWIM11, 5 – 14.
- Downing, R.A. (1972) A summary of saline intrusion problems in the United Kingdom, SWIM3, 70 – 74.

- Flathe, N. (1968) Stand der Kenntnisse über den Anwendungsbereich der Geolektrik im niedersächsischen Küstengebiet (State of knowledge on the scope of geoelectics in the coastal area of Lower Saxony). SWIM1, 6 – 9.
- Geyh, M.A. (1968) <sup>14</sup>C- und <sup>3</sup>H-Konzentrationsbestimmungen (<sup>14</sup>C and <sup>3</sup>H concentration determinations), SWIM1, 16 – 19.
- Højberg, A. L., Hansen, A. L., Wachniew, P., Żurek, A. J., Virtanen, S., Arustiene, J., Refsgaard, J. C. (2017). Review and assessment of nitrate reduction in groundwater in the Baltic Sea Basin. *Journal of Hydrology: Regional Studies*, 12, 50-68.
- Klinge, H., Boehme, J., Ludwig, R. (1999) Fresh-water/salt-water distribution in the aquifer system above the Gorleben salt dome: Results of the Gorleben site investigation programme. SWIM 15, 172-177
- Lebbe, L. (1981) The subterranean flow of fresh and salt water underneath the western Belgian beach, SWIM7, 193 – 219.
- Lindewald, H. (1981) Saline groundwater in Sweden. SWIM7, 24 – 32.
- Michael, H.A., Post, V.E.A., Wilson, A.M., Werner, A.D., 2017. Science, society, and the coastal groundwater squeeze. *Water Resources Research*, 53(4): 2610-2617. DOI:10.1002/2017wr020851
- Nielsen, H. (1968) Schwefelisotopen-Untersuchungen (Sulfur isotope studies), SWIM1, 13 – 14.
- Ochmann, N., Fielitz, K. (1993) Estimation of horizontal and vertical groundwater flow from well logging and pressure data in groundwater of variable density above a salt dome. SWIM12, 359-369
- Scholten, J., Kreuzburg, M., Knoeller, K., Rapaglia, J., Schlüter, M., Schubert, M. (2014) Submarine groundwater discharge in the southwestern Baltic Sea. SWIM23, 362.
- Schwerdtfeger, B. (1981) On the occurrence of submarine fresh-water discharges. *Geologisches Jahrbuch Reihe C* 29, 231 – 240.
- Stahl, W. (1968) Sauerstoffisotopen-Untersuchungen (Oxygen isotopes - Investigations), SWIM1, 14 – 16.
- Sund, B. and G. Bergman (1981) Sea water intrusion in drilled wells, SWIM7, 45 – 58.
- SWIM (2018) Salt Water Intrusion Meeting (SWIM). <http://www.swim-site.org>. Accessed Jan 2018

Szymczycha, B., & Pempkowiak, J. (2016). Research on Submarine Groundwater Discharge in the Baltic Sea. In: *The Role of Submarine Groundwater Discharge as Material Source to the Baltic Sea* (pp. 53-132). Springer International Publishing.

Verruijt, A. (1972) Analysis of interface problems by the finite element method. *SWIM3*, 53 – 59.

Zechner et al. (2014) Simulation of high-contrast density driven transport. *SWIM22*, 247-250.

Zuber, A., Kozerski, B., Sadurski, A., Kwaterkiewicz, A., Grabczak, J. (1990) Origin of brackish waters in the Quaternary aquifer of the Vistula delta. *SWIM11*, 249 – 262.




FIGURE CAPTION:

**Figure 1** Maps with dots showing the locations of the case studies of saltwater intrusion presented at SWIMs. The main map covers the entire globe, while the inset map is an enlargement of Europe and the Mediterranean region, which have the highest density of studies. When the precise location of a study could not be determined, the dot was placed in the centre of a country.

